

COTENSOR COALGEBRAS IN MONOIDAL CATEGORIES

A. ARDIZZONI, C. MENINI AND D. ŞTEFAN

ABSTRACT. We introduce the concept of cotensor coalgebra for a given bicomodule over a coalgebra in an abelian monoidal category. Under some further conditions we show that such a cotensor coalgebra exists and satisfies a meaningful universal property. We prove that this coalgebra is formally smooth whenever the comodule is relative injective and the coalgebra itself is formally smooth.

INTRODUCTION

Let C be a coalgebra over a field k and let M be a C -bicomodule. The cotensor coalgebra $T_C^c(M)$ was introduced by Nichols in [Ni] as a main tool to construct some new Hopf algebras that he called "bialgebras of type one". These bialgebras can be reconstructed, via a bosonization procedure, from the so called Nichols algebras, which are essentially the H -coinvariant elements of the bialgebras of type one, in the case when $C = H$ is a Hopf algebra and M is a Hopf bimodule. Nichols algebras, also named quantum symmetric algebras in [Ro], have been deeply investigated and appear as a main step in the classification of finite dimensional Hopf algebras problem (see, e.g., [AG] and [AS]). In fact, in the case that $C = H$ is a Hopf algebra and M is a Hopf H -bimodule, the cotensor coalgebra $T_C^c(M)$ is a bialgebra that is called "quantum shuffle Hopf algebra" by Rosso in [Ro] where some fundamental properties of this bialgebra and of its coinvariant Hopf algebra are investigated. The coalgebra of paths of a quiver Q is an instance of a cotensor coalgebra. Namely let Q_0 be the set of vertices and let Q_1 be the set of arrows of Q . Then $M = KQ_1$ is a C -bicomodule where $C = KQ_0$ is equipped with its natural coalgebra structure. The cotensor coalgebra $T_C^c(M)$ is the path coalgebra of the quiver Q . In [CR], Cibils and Rosso provide the classification of path coalgebras which admit a graded Hopf algebra structure, allowing the quiver to be infinite. On the other hand, in [JLMS], hereditary coalgebras with coseparable coradical are characterized by means of a suitable cotensor coalgebra. Moreover it is proved that if C is a formally smooth coalgebra (see Definition 4.12) and M is $\mathcal{I}$ -injective (see 4.2) then $T_C^c(M)$ is formally smooth.

In this paper we introduce the notion of cotensor coalgebra in an abelian monoidal category. We would like to outline that this fact is not immediate. In fact the notion of coradical plays a fundamental role in the usual definition for coalgebras over a field (see [Ni]) while we have no coradical substitution here. Also, having developed in [AMS] the notion of formally smooth coalgebras for abelian monoidal categories, we wanted to obtain the second quoted result of [JLMS] in this more general setting, namely we prove Theorem 4.11. In a forthcoming paper we will deal with the other quoted result, establishing a suitable criterion.

The paper is organized as follows.

In Section 1 we give the definition of abelian monoidal category (see Definition 1.4) and prove some preliminary results on direct limits of direct systems in monoidal categories.

Section 2 is devoted to the construction of the cotensor coalgebra. Its coalgebra structure is obtained (see Theorem 2.9) as a direct limit of a direct system of certain coalgebras defined by means of Hochschild 2-cocycles. In Theorem 2.15 we show that such a cotensor coalgebra exists and satisfies a meaningful universal property.

1991 *Mathematics Subject Classification.* Primary 18D10; Secondary 18A30.

Key words and phrases. Monoidal categories, colimits, wedge products, Cotensor Coalgebras.

This paper was written while A. Ardizzoni and C. Menini were members of G.N.S.A.G.A. with partial financial support from M.I.U.R..

Section 3 deals with some technical results that we will use in Section 4 where we treat formal smoothness of the cotensor coalgebra (see Theorem 4.15).

Finally, in Section 5, we relate our results with the classical one of vector spaces and prove a criterion (Theorem 5.7) which establishes a strong universal property of the cotensor coalgebra for the main abelian monoidal categories related to a Hopf algebra.

Notation. In a category $\mathcal{M}$ the set of morphisms from X to Y will be denoted by $\mathcal{M}(X, Y)$. If X is an object in $\mathcal{M}$ then the functor $\mathcal{M}(X, -)$ from $\mathcal{M}$ to $\mathbf{Sets}$ associates to any morphism $u : U \rightarrow V$ in $\mathcal{M}$ the function that will be denoted by $\mathcal{M}(X, u)$.

1. DIRECT LIMITS IN MONOIDAL CATEGORIES

1.1. A *monoidal category* means a category $\mathcal{M}$ that is endowed with a functor $\otimes : \mathcal{M} \times \mathcal{M} \rightarrow \mathcal{M}$, an object $\mathbf{1} \in \mathcal{M}$ and functorial isomorphisms: $a_{X,Y,Z} : (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$, $l_X : \mathbf{1} \otimes X \rightarrow X$ and $r_X : X \otimes \mathbf{1} \rightarrow X$. The functorial morphism a is called the *associativity constraint* and satisfies the *Pentagon Axiom*, that is the following diagram

$$\begin{array}{ccccc}
 & & (U \otimes V) \otimes (W \otimes X) & & \\
 & \nearrow a_{U \otimes V, W, X} & & \nwarrow a_{U, V, W \otimes X} & \\
 ((U \otimes V) \otimes W) \otimes X & & & & U \otimes (V \otimes (W \otimes X)) \\
 \searrow a_{U, V, W \otimes X} & & & & \nearrow U \otimes a_{V, W, X} \\
 (U \otimes (V \otimes W)) \otimes X & \xrightarrow{a_{U, V \otimes W, X}} & U \otimes ((V \otimes W) \otimes X) & &
 \end{array}$$

is commutative, for every U, V, W, X in $\mathcal{M}$. The morphisms l and r are called the *unit constraints* and they are assumed to satisfy the *Triangle Axiom*, i.e. the following diagram

$$\begin{array}{ccc}
 (V \otimes \mathbf{1}) \otimes W & \xrightarrow{a_{V, \mathbf{1}, W}} & V \otimes (\mathbf{1} \otimes W) \\
 \searrow r_V \oplus W & & \swarrow V \oplus l_W \\
 & V \otimes W &
 \end{array}$$

is commutative. The object $\mathbf{1}$ is called the *unit* of $\mathcal{M}$. For details on monoidal categories we refer to [Ka, Chapter XI] and [Maj]. A monoidal category is called *strict* if the associativity constraint and unit constraints are the corresponding identity morphisms.

1.2. As it is noticed in [Maj, p. 420], the Pentagon Axiom solves the consistency problem that appears because there are two ways to go from $((U \otimes V) \otimes W) \otimes X$ to $U \otimes (V \otimes (W \otimes X))$. The coherence theorem, due to S. Mac Lane, solves the similar problem for the tensor product of an arbitrary number of objects in $\mathcal{M}$. Accordingly with this theorem, we can always omit all brackets and simply write $X_1 \otimes \cdots \otimes X_n$ for any object obtained from $X_1, \dots, X_n$ by using $\otimes$ and brackets. Also as a consequence of the coherence theorem, the morphisms a, l, r take care of themselves, so they can be omitted in any computation involving morphisms in $\mathcal{M}$. The notions of algebra, module over an algebra, coalgebra and comodule over a coalgebra can be introduced in the general setting of monoidal categories. For more details, see [AMS].

PROPOSITION 1.3. *Let $\mathcal{M}$ be a monoidal category. Assume that $\mathcal{M}$ is also an abelian category. Let C and D be coalgebras in $\mathcal{M}$. Let $X, Y \in {}^C\mathcal{M}^D$ and let $f : X \rightarrow Y$ be a morphism of (C, D) -bicomodules. Let $\mathbb{H} : {}^C\mathcal{M}^D \rightarrow \mathcal{M}$ be the forgetful functor. Then*

1) *Coker($\mathbb{H}(f)$) carries a natural (C, D) -bicomodule structure (such that the definition map is a morphism of bicomodules) that makes it the cokernel of f in ${}^C\mathcal{M}^D$.*

2) if the functor $C \otimes (-) \otimes D : \mathcal{M} \rightarrow \mathcal{M}$ is left exact (e.g. the tensor functors are left exact), then $\text{Ker}(\mathbb{H}(f))$ carries a natural (C, D) -bicomodule structure (such that the definition map is a morphism of bicomodules) that makes it the kernel of f in ${}^C\mathcal{M}^D$.

Proof. Since the dual category of an abelian category is an abelian category, the conclusion follows by applying [Ar2, Proposition 3.3]. $\square$

The previous proposition justifies the following definition.

DEFINITION 1.4. A monoidal category $(\mathcal{M}, \otimes, \mathbf{1})$ will be called an **abelian monoidal category** if:

- (1) $\mathcal{M}$ is an abelian category
- (2) both the functors $X \otimes (-) : \mathcal{M} \rightarrow \mathcal{M}$ and $(-) \otimes X : \mathcal{M} \rightarrow \mathcal{M}$ are additive and left exact, for every object $X \in \mathcal{M}$.

REMARK 1.5. The above definition of abelian monoidal category is dual to [AMS, Definition 1.8], where algebras were investigated. Therefore, we should have used the terminology "dual abelian monoidal category", which we had not for matter of style.

1.6. Let E be a coalgebra in an abelian monoidal category $\mathcal{M}$. Let us recall, (see [Mo, page 60]), the definition of wedge product of two subobjects X, Y of E in $\mathcal{M}$:

$$X \wedge_E Y := \text{Ker}[(p_X \otimes p_Y) \circ \Delta_E],$$

where $p_X : E \rightarrow E/X$ and $p_Y : E \rightarrow E/Y$ are the canonical quotient maps.

PROPOSITION 1.7. Let $(\mathcal{M}, \otimes, \mathbf{1})$ be an abelian monoidal category. Let (C, Δ, ε) be a coalgebra in $\mathcal{M}$ and let L be a C -bicomodule. Let $f : C \rightarrow L$ be a morphism in ${}^C\mathcal{M}^C$, where C is regarded as a bicomodule via Δ . Then $(D, \delta) := \text{Ker}(f)$ carries a natural coalgebra structure such that δ is a morphism of coalgebras.

Proof. By Proposition 1.3, D is a C -bicomodule and δ is a morphism of bicomodules. Denote by ρ_D^l and ρ_D^r the left and the right C -comodule structure of D respectively. By left exactness of the tensor functors, we have that $(D \otimes D, D \otimes \delta) = \text{Ker}(D \otimes f)$. Consider the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & D \otimes D & \xrightarrow{D \otimes \delta} & D \otimes C & \xrightarrow{D \otimes f} & D \otimes L \\ & & \searrow \Delta_D & & \uparrow \rho_D^r & & \\ & & & & D & & \end{array}$$

We have:

$$(\delta \otimes L)(D \otimes f)\rho_D^r = (C \otimes f)(\delta \otimes C)\rho_D^r = (C \otimes f)\Delta\delta = \rho_L^l f\delta = 0.$$

By left exactness of the tensor functors, $\delta \otimes L$ is a monomorphism, so that we get $(D \otimes f)\rho_D^r = 0$ and hence, by the universal property of the kernel, there exists a unique morphism $\Delta_D : D \rightarrow D \otimes D$ in $\mathcal{M}$ such that $(D \otimes \delta)\Delta_D = \rho_D^r$.

Let us prove that Δ_D is coassociative. Since

$$(1) \quad (\delta \otimes \delta)\Delta_D = (\delta \otimes C)(D \otimes \delta)\Delta_D = (\delta \otimes C)\rho_D^r = \Delta\delta,$$

we get:

$$\begin{aligned} (\delta \otimes \delta \otimes \delta)(\Delta_D \otimes D)\Delta_D &= (\Delta \otimes C)(\delta \otimes \delta)\Delta_D \\ &= (\Delta \otimes C)\Delta\delta \\ &= (C \otimes \Delta)\Delta\delta \\ &= (C \otimes \Delta)(\delta \otimes \delta)\Delta_D = (\delta \otimes \delta \otimes \delta)(D \otimes \Delta_D)\Delta_D. \end{aligned}$$

By left exactness of the tensor functors, $\delta \otimes \delta \otimes \delta = (C \otimes C \otimes \delta)(\delta \otimes C \otimes D)(D \otimes \delta \otimes D)$ is a monomorphism so that we obtain $(\Delta_D \otimes D)\Delta_D = (D \otimes \Delta_D)\Delta_D$.

Set $\varepsilon_D := \varepsilon_C \delta : D \rightarrow \mathbf{1}$. Then we have:

$$(\delta \otimes \mathbf{1})(D \otimes \varepsilon_D)\Delta_D = (C \otimes \varepsilon_C)(\delta \otimes \delta)\Delta_D = (C \otimes \varepsilon_C)\Delta\delta = r_C^{-1}\delta = (\delta \otimes \mathbf{1})r_D^{-1}.$$

Since $(\delta \otimes \mathbf{1})$ is a monomorphism, we have $(D \otimes \varepsilon_D)\Delta_D = r_D^{-1}$. Analogously one gets $(\varepsilon_D \otimes D)\Delta_D = l_D^{-1}$. Thus $(D, \Delta_D, \varepsilon_D)$ is a coalgebra and by relation (1) and definition of ε_D , δ is a homomorphism of coalgebras. $\square$

PROPOSITION 1.8. *Let $\mathcal{M}$ be a monoidal category with direct limits. Let $((X_i)_{i \in \mathbb{N}}, (\xi_i^j)_{i,j \in \mathbb{N}})$ be a direct system in $\mathcal{M}$, where, for $i \leq j$, $\xi_i^j : X_i \rightarrow X_j$. Assume that X_i is a coalgebra and that ξ_i^j is a homomorphism of coalgebras for any $i, j \in \mathbb{N}$. Then $\varinjlim X_i$ carries a natural coalgebra structure that makes it the direct limit of $((X_i)_{i \in \mathbb{N}}, (\xi_i^j)_{i,j \in \mathbb{N}})$ as a direct system of coalgebras.*

Proof. Let $(X_i, \Delta_{X_i}, \varepsilon_{X_i})$ be a coalgebra in $(\mathcal{M}, \otimes, \mathbf{1})$ for any $i \in \mathbb{N}$. Set $X := \varinjlim X_i$. Let $(\xi_i : X_i \rightarrow X)_{i \in \mathbb{N}}$ be the structural morphism of the direct limit, so that $\xi_j \xi_i^j = \xi_i$ for any $i \leq j$. We put

$$\Delta_i = (\xi_i \otimes \xi_i)\Delta_{X_i} : X_i \rightarrow X \otimes X, \text{ for any } i \in \mathbb{N}.$$

Since ξ_i^j is a homomorphism of coalgebras, we have that:

$$\Delta_j \xi_i^j = (\xi_j \otimes \xi_j)\Delta_{X_j} \xi_i^j = (\xi_j \otimes \xi_j)(\xi_i^j \otimes \xi_i^j)\Delta_{X_i} = (\xi_i \otimes \xi_i)\Delta_{X_i} = \Delta_i,$$

so that there exists a unique morphism $\Delta : X \rightarrow X \otimes X$ such that

$$(2) \quad \Delta \xi_i = \Delta_i = (\xi_i \otimes \xi_i)\Delta_{X_i} \text{ for any } i \in \mathbb{N}.$$

Let us prove that Δ is coassociative:

$$\begin{aligned} (X \otimes \Delta)\Delta \xi_i &= (X \otimes \Delta)(\xi_i \otimes \xi_i)\Delta_{X_i} \\ &= (\xi_i \otimes \xi_i \otimes \xi_i)(X_i \otimes \Delta_{X_i})\Delta_{X_i} \\ &= (\xi_i \otimes \xi_i \otimes \xi_i)(\Delta_{X_i} \otimes X_i)\Delta_{X_i} \\ &= (\Delta \otimes X)(\xi_i \otimes \xi_i)\Delta_{X_i} = (\Delta \otimes X)\Delta \xi_i. \end{aligned}$$

Since the relation above holds true for any $i \in \mathbb{N}$, by the universal property of the direct limit we deduce that $(X \otimes \Delta)\Delta = (\Delta \otimes X)\Delta$. Now, as ξ_i^j is a homomorphism of coalgebras, $\varepsilon_{X_j} \xi_i^j = \varepsilon_{X_i}$. Hence, there exists a unique morphism $\varepsilon : X \rightarrow \mathbf{1}$ such that

$$(3) \quad \varepsilon \xi_i = \varepsilon_{X_i} \text{ for any } i \in \mathbb{N}.$$

Then we have:

$$(X \otimes \varepsilon)\Delta \xi_i = (X \otimes \varepsilon)(\xi_i \otimes \xi_i)\Delta_{X_i} = (\xi_i \otimes \mathbf{1})(X_i \otimes \varepsilon_{X_i})\Delta_{X_i} = (\xi_i \otimes \mathbf{1})r_{X_i}^{-1} = r_X^{-1}\xi_i.$$

Since the relation above holds true for any $i \in \mathbb{N}$, by the universal property of direct limits we deduce that $(X \otimes \varepsilon)\Delta = r_X^{-1}$. Analogously one gets $(\varepsilon \otimes X)\Delta = l_X^{-1}$. Thus (X, Δ, ε) is a coalgebra in $\mathcal{M}$. Note that relations (2) and (3) mean that $\xi_i : X_i \rightarrow X$ is a homomorphism of coalgebras. Let now $(C, \Delta_C, \varepsilon_C)$ be a coalgebra in $\mathcal{M}$ and let $(f_i : X_i \rightarrow C)_{i \in \mathbb{N}}$ be a compatible family of morphisms of coalgebras in $\mathcal{M}$. Since $(f_i : X_i \rightarrow C)_{i \in \mathbb{N}}$ is a compatible family of morphisms in $\mathcal{M}$, there exists a unique morphism $f : X \rightarrow C$ such that $f \xi_i = f_i$ for any $i \in \mathbb{N}$. We prove that f is a homomorphism of coalgebras. We have:

$$(f \otimes f)\Delta \xi_i = (f \otimes f)(\xi_i \otimes \xi_i)\Delta_{X_i} = (f_i \otimes f_i)\Delta_{X_i} = \Delta_C f_i = \Delta_C f \xi_i; \quad \varepsilon_C f \xi_i = \varepsilon_C f_i = \varepsilon_{X_i} = \varepsilon \xi_i.$$

Since the relations above hold true for any $i \in \mathbb{N}$, by the universal property of the direct limit we deduce that

$$(f \otimes f)\Delta = \Delta_C f; \quad \varepsilon_C f = \varepsilon.$$

$\square$

1.9. Let X be an object in an abelian monoidal category $(\mathcal{M}, \otimes, \mathbf{1})$. Set

$$X^{\otimes 0} = \mathbf{1}, \quad X^{\otimes 1} = X \quad \text{and} \quad X^{\otimes n} = X^{\otimes n-1} \otimes X, \text{ for every } n > 1$$

and for every morphism $f : X \rightarrow Y$ in $\mathcal{M}$, set

$$f^{\otimes 0} = \text{Id}_{\mathbf{1}}, \quad f^{\otimes 1} = f \quad \text{and} \quad f^{\otimes n} = f^{\otimes n-1} \otimes f, \text{ for every } n > 1.$$

Let $(C, \Delta_C, \varepsilon_C)$ be a coalgebra in $\mathcal{M}$ and for every $n \in \mathbb{N}$, define the n^{th} iterated comultiplication of C , $\Delta_C^n : C \rightarrow C^{\otimes n+1}$, by

$$\Delta_C^0 = \text{Id}_C, \quad \Delta_C^1 = \Delta_C \quad \text{and} \quad \Delta_C^n = (\Delta_C^{\otimes n-1} \otimes C) \Delta_C, \text{ for every } n > 1.$$

Let $\delta : D \rightarrow C$ be a monomorphism which is a homomorphism of coalgebras in $\mathcal{M}$. Denote by (L, p) the cokernel of δ in $\mathcal{M}$. Regard D as a C -bicomodule via δ and observe that, by Proposition 1.3, L is a C -bicomodule and p is a morphism of bicomodules. Let

$$(D^{\wedge_C^n}, \delta_n) := \text{Ker}(p^{\otimes n} \Delta_C^{n-1})$$

for any $n \in \mathbb{N} \setminus \{0\}$. Note that $(D^{\wedge_C^1}, \delta_1) = (D, \delta)$ and $(D^{\wedge_C^2}, \delta_2) = D \wedge_C D$.

In order to simplify the notations we set $(D^{\wedge_C^0}, \delta_0) = (0, 0)$.

Now, if we assume that $\mathcal{M}$ has left exact tensor functors, by Proposition 1.7, since $p^{\otimes n} \Delta_C^{n-1}$ is a morphism of C -bicomodules (as a composition of morphisms of C -bicomodules), we get that $D^{\wedge_C^n}$ is a coalgebra and $\delta_n : D^{\wedge_C^n} \rightarrow C$ is a coalgebra homomorphism for any $n > 0$ and hence for any $n \in \mathbb{N}$.

PROPOSITION 1.10. *Let $\delta : D \rightarrow C$ be a monomorphism which is a homomorphism of coalgebras in an abelian monoidal category $\mathcal{M}$. Then for any $i \leq j$ in $\mathbb{N}$ there is a (unique) morphism $\xi_i^j : D^{\wedge_C^i} \rightarrow D^{\wedge_C^j}$ such that*

$$(4) \quad \delta_j \xi_i^j = \delta_i.$$

Moreover ξ_i^j is a coalgebra homomorphism and $((D^{\wedge_C^i})_{i \in \mathbb{N}}, (\xi_i^j)_{i, j \in \mathbb{N}})$ is a direct system in $\mathcal{M}$ whose direct limit, if it exists, carries a natural coalgebra structure that makes it the direct limit of $((D^{\wedge_C^i})_{i \in \mathbb{N}}, (\xi_i^j)_{i, j \in \mathbb{N}})$ as a direct system of coalgebras.

Proof. Set $D^i := D^{\wedge_C^i}$ for any $i \in \mathbb{N}$. Consider the following diagram:

$$\begin{array}{ccccc} 0 & \longrightarrow & D^{i+1} & \xrightarrow{\delta_{i+1}} & C & \xrightarrow{p^{\otimes i+1} \Delta_C^i} & L^{\otimes i+1} \\ & & & \nwarrow \xi_i^{i+1} & \uparrow \delta_i & & \\ & & & & D^i & & \end{array}$$

Let $i > 0$. Since $\delta_i = \text{Ker}(p^{\otimes i} \Delta_C^{i-1})$ is a coalgebra homomorphism, we have:

$$\begin{aligned} p^{\otimes i+1} \Delta_C^i \delta_i &= p^{\otimes i+1} (C \otimes \Delta_C^{i-1}) \Delta_C \delta_i \\ &= p^{\otimes i+1} (C \otimes \Delta_C^{i-1}) (\delta_i \otimes \delta_i) \Delta_{D^i} = (p \delta_i \otimes p^{\otimes i} \Delta_C^{i-1} \delta_i) \Delta_{D^i} = 0. \end{aligned}$$

Then, for any $i \geq 1$, by the universal property of the kernel, there exists a unique morphism $\xi_i^{i+1} : D^i \rightarrow D^{i+1}$ such that $\delta_{i+1} \xi_i^{i+1} = \delta_i$. Set $\xi_0^1 = 0$ and for any $j > i$, define:

$$\xi_i^j = \xi_{j-1}^j \xi_{j-2}^{j-1} \dots \xi_{i+1}^{i+2} \xi_i^{i+1} : D^i \rightarrow D^j.$$

In such a way we obviously obtain a direct system in $\mathcal{M}$. Let us prove that ξ_i^j is a homomorphism of coalgebras for any $j > i$. It is clearly sufficient to verify this for $j = i + 1$.

As δ_{i+1} and δ_i are coalgebra homomorphisms, we have

$$(\delta_{i+1} \otimes \delta_{i+1}) \Delta_{D^{i+1}} \xi_i^{i+1} = \Delta_D \delta_{i+1} \xi_i^{i+1} = \Delta_D \delta_i = (\delta_i \otimes \delta_i) \Delta_{D^i} = (\delta_{i+1} \otimes \delta_{i+1}) (\xi_i^{i+1} \otimes \xi_i^{i+1}) \Delta_{D^i}.$$

Since the tensor functors are left exact, $\delta_{i+1} \otimes \delta_{i+1}$ is a monomorphism so that we get $\Delta_{D^{i+1}} \xi_i^{i+1} = (\xi_i^{i+1} \otimes \xi_i^{i+1}) \Delta_{D^i}$. Moreover we have

$$\varepsilon_{D^{i+1}} \xi_i^{i+1} = \varepsilon_D \delta_{i+1} \xi_i^{i+1} = \varepsilon_D \delta_i = \varepsilon_{D^i}.$$

The last assertion follows by Proposition 1.8. $\square$

NOTATION 1.11. Let $\delta : D \rightarrow C$ be a homomorphism of coalgebras in a cocomplete abelian monoidal category $\mathcal{M}$. By Proposition 1.10 $((D^{\wedge^i_C})_{i \in \mathbb{N}}, (\xi_i^j)_{i,j \in \mathbb{N}})$ is a direct system in $\mathcal{M}$ whose direct limit carries a natural coalgebra structure that makes it the direct limit of $((D^{\wedge^i_C})_{i \in \mathbb{N}}, (\xi_i^j)_{i,j \in \mathbb{N}})$ as a direct system of coalgebras.

From now on we set: $(\tilde{D}_C, (\xi_i)_{i \in \mathbb{N}}) = \varinjlim (D^{\wedge^i_C})_{i \in \mathbb{N}}$, where $\xi_i : D^{\wedge^i_C} \rightarrow \tilde{D}_C$ denotes the structural morphism of the direct limit. We simply write $\tilde{D}$ if there is no danger of confusion. We note that, since $\tilde{D}$ is a direct limit of coalgebras, the canonical (coalgebra) homomorphisms $(\delta_i : D^{\wedge^i_C} \rightarrow C)_{i \in \mathbb{N}}$, which are compatible by (4), factorize to a unique coalgebra homomorphism $\tilde{\delta} : \tilde{D} \rightarrow C$ such that $\tilde{\delta}\xi_i = \delta_i$ for any $i \in \mathbb{N}$.

2. COTENSOR COALGEBRA

2.1. Let (C, Δ, ε) be an coalgebra in $(\mathcal{M}, \otimes, \mathbf{1})$ and let (L, ρ_L^l, ρ_L^r) be a C -bicomodule. Recall that a morphism $\zeta : L \rightarrow C \otimes C$ is called a Hochschild 2-cocyle whenever

$$b^2(\zeta) = (\zeta \otimes C) \circ \rho_L^r - (C \otimes \Delta) \circ \zeta + (\Delta \otimes C) \circ \zeta - (C \otimes \zeta) \circ \rho_L^l$$

is zero. See 4.1.

DEFINITION 2.2. Let $(C, \Delta_C, \varepsilon_C)$ be a coalgebra in $(\mathcal{L}, \otimes, \mathbf{1})$ and let (L, ρ_l, ρ_r) be a bicomodule over C . A *Hochschild extension* of C with cokernel L , is an exact sequence in $\mathcal{M}$:

$$(E) \quad 0 \longrightarrow C \xrightarrow{\sigma} E \xrightarrow{p} L \longrightarrow 0$$

that satisfy the following conditions:

- a) $(E, \Delta_E, \varepsilon_E)$ is a coalgebra in $\mathcal{M}$;
- b) σ is a homomorphism of coalgebras that has a retraction π in $\mathcal{M}$;
- c) $C \wedge_E C = E$, that is $(p \otimes p)\Delta = 0$;
- d) the morphisms ρ_l and ρ_r fulfill the following relations

$$\rho_l p = (\pi \otimes p)\Delta_E \quad \rho_r p = (p \otimes \pi)\Delta_E.$$

The following result will lead to the definition of a coalgebra structure for the cotensor coalgebra.

LEMMA 2.3. Let (C, Δ, ε) be a coalgebra in an abelian monoidal category $\mathcal{M}$ and let (L, ρ_L^r, ρ_L^l) a C -bicomodule. Suppose that $\zeta : L \rightarrow C \otimes C$ is a morphism in $\mathcal{M}$. Define $\Delta_\zeta : C \oplus L \rightarrow (C \oplus L) \otimes (C \oplus L)$ and $\varepsilon_\zeta : C \oplus L \rightarrow \mathbf{1}$ by:

$$(5) \quad \Delta_\zeta := (i_C \otimes i_C) \Delta p_C + [(i_L \otimes i_C) \rho_L^r + (i_C \otimes i_L) \rho_L^l - (i_C \otimes i_C) \zeta] p_L,$$

$$(6) \quad \varepsilon_\zeta := \varepsilon p_C + l_1 (\varepsilon \otimes \varepsilon) \zeta p_L,$$

where i_C, i_L are the canonical injections in $C \oplus L$ and p_C, p_L are the canonical projections. Then Δ_ζ is a coassociative comultiplication if and only if ζ is a Hochschild 2-cocycle. Moreover, in this case, $(C \oplus L, \Delta_\zeta, \varepsilon_\zeta)$ is a coalgebra and $i_C : C \rightarrow (C \oplus L, \Delta_\zeta, \varepsilon_\zeta)$ is a Hochschild extension of C with cokernel (L, p_L) . This extension will be denoted by E_ζ .

Proof. The dual of an abelian monoidal category is an abelian monoidal category in the sense of [AMS, Definition 1.8]. The conclusion follows by applying [AMS, Lemma 2.5]. $\square$

Let $\mathcal{M}$ be an abelian monoidal category and let C be a coalgebra in $(\mathcal{M}, \otimes, \mathbf{1})$. Given a right C -bicomodule (V, ρ_V^r) and a left C -comodule (W, ρ_W^l) , their cotensor product over C in $\mathcal{M}$ is defined to be the equalizer $(V \square_C W, \chi)$ of the couple of morphism $(\rho_V^r \otimes W, V \otimes \rho_W^l)$:

$$0 \longrightarrow V \square_C W \xrightarrow{\chi} V \otimes W \xrightleftharpoons[V \otimes \rho_W^l]{\rho_V^r \otimes W} V \otimes C \otimes W.$$

Since the tensor functors are left exact, in view of Proposition 1.3, then $V \square_C W$ is also a C -bicomodule, namely it is C -sub-bicomodule of $V \otimes W$, whenever V and W are C -bicomodules. Furthermore, in this case, the category $({}^C\mathcal{M}^C, \square_C, C)$ is still an abelian monoidal category; the associative and unit constraints are induced by the ones in $\mathcal{M}$ (the proof is dual to [AMS, Theorem

1.11]). Therefore, also using $\square_C$, one can forget about brackets. Moreover the functors $M\square_C(-) : {}^C\mathcal{M} \rightarrow \mathcal{M}$ and $(-)\square_C M : \mathcal{M}^C \rightarrow \mathcal{M}$ are left exact for any $M \in \mathcal{M}$.

We will write $\square$ instead of $\square_C$, whenever there is no danger of misunderstanding.

2.4. Let (C, Δ, ε) be a coalgebra in a cocomplete abelian monoidal category $\mathcal{M}$ and let (M, ρ_M^r, ρ_M^l) be a C -bicomodule. Set

$$M^{\square 0} = C, M^{\square 1} = M \quad \text{and} \quad M^{\square n} = M^{\square n-1} \square M \text{ for any } n > 1$$

and define $(C^n(M))_{n \in \mathbb{N}}$ by

$$C^0(M) = 0, C^1(M) = C \quad \text{and} \quad C^n(M) = C^{n-1}(M) \oplus M^{\square n-1} \text{ for any } n > 1.$$

Let $\sigma_i^{i+1} : C^i(M) \rightarrow C^{i+1}(M)$ be the canonical inclusion and for any $j > i$, define:

$$\sigma_i^j = \sigma_{j-1}^j \sigma_{j-2}^{j-1} \cdots \sigma_{i+1}^{i+2} \sigma_i^{i+1} : C^i(M) \rightarrow C^j(M).$$

Then $((C^i(M))_{i \in \mathbb{N}}, (\sigma_i^j)_{i,j \in \mathbb{N}})$ is a direct system in $\mathcal{M}$. We set

$$T_C^c(M) = \bigoplus_{n \in \mathbb{N}} M^{\square n} = C \oplus M \oplus M^{\square 2} \oplus M^{\square 3} \oplus \cdots$$

and we denote by $\sigma_i : C^i(M) \rightarrow T_C^c(M)$ the canonical inclusion.

Throughout let

$$\begin{aligned} \pi_n^m &: C^n(M) \rightarrow C^m(M) \quad (m \leq n), & \pi_n &: T_C^c(M) \rightarrow C^n(M), \\ p_n^m &: C^m(M) \rightarrow M^{\square m} \quad (m < n), & p_n &: T_C^c(M) \rightarrow M^{\square n}, \end{aligned}$$

be the canonical projections and let

$$\begin{aligned} \sigma_n^m &: C^m(M) \rightarrow C^n(M) \quad (m \leq n), & \sigma_n &: C^n(M) \rightarrow T_C^c(M), \\ i_m^n &: M^{\square m} \rightarrow C^n(M) \quad (m < n), & i_m &: M^{\square m} \rightarrow T_C^c(M), \end{aligned}$$

be the canonical injection for any $m, n \in \mathbb{N}$.

For technical reasons we set $\pi_n^m = 0, \sigma_n^m = 0$ for any $n < m$ and $p_n^m = 0, i_m^n = 0$ for any $n \leq m$. Then, we have the following relations:

$$p_n \sigma_k = p_k^n, \quad p_n i_k = \delta_{n,k} \text{Id}_{M^{\square k}}, \quad \pi_n i_k = i_k^n.$$

Moreover, we have:

$$\begin{aligned} \pi_n^m \sigma_k^n &= \sigma_k^m, \text{ if } k \leq m \leq n, & \text{and} & & \pi_n^m \sigma_k^n &= \pi_k^m, \text{ if } m \leq k \leq n, \\ p_n^m \pi_k^n &= p_k^m, \text{ if } m < n \leq k, & \text{and} & & \sigma_n^m i_k^n &= i_k^m, \text{ if } k < n \leq m, \\ p_n^m \sigma_k^n &= p_k^m, \text{ if } m < k \leq n, & \text{and} & & \pi_n^m i_k^n &= i_k^m, \text{ if } k < m \leq n, \\ p_n^m \pi_n &= p_n, \text{ if } m < n, & \text{and} & & \sigma_n i_m^n &= i_m, \text{ if } m < n, \\ \pi_n \sigma_k &= \sigma_k^n, \text{ if } k \leq n, & \text{and} & & \pi_n \sigma_k &= \pi_k^n, \text{ if } n \leq k, \\ p_n^m i_m^n &= \text{Id}_{M^{\square m}}, \text{ if } m < n. \end{aligned}$$

In the other cases, these compositions are zero.

PROPOSITION 2.5. *Let $\mathcal{M}$ be a cocomplete abelian category. Let $(V_i)_{i \in \mathbb{N}}$ be a family of objects in $\mathcal{M}$ and let $(V, v_i) = \bigoplus_{i \in \mathbb{N}} V_i$ be the direct sum of the family $(V_i)_{i \in \mathbb{N}}$. Then*

$$(V, \nabla_{i=0}^n(v_i)) = \varinjlim (\bigoplus_{i=0}^n V_i),$$

where $\nabla_{i=0}^n(v_i) : \bigoplus_{i=0}^n V_i \rightarrow V$ denotes the codiagonal morphism associated to the family $(v_i)_{i=0}^n$.

Proof. Set $V^n := \bigoplus_{i=0}^n V_i$, for any $n \in \mathbb{N}$, and let $w_m^n : V^m \rightarrow V^n$ be the canonical inclusion for $m \leq n$. Let $(f_n : V^n \rightarrow X)_n$ be a compatible family of morphisms in $\mathcal{M}$, i.e. $f_n w_m^n = f_m$ for any $m \leq n$. Let $v_m^n : V_m \rightarrow V^n$ be the canonical inclusion for every $m \leq n$ and let $v_m^n = 0$ otherwise. Note that the morphism $\nabla_{i=0}^n(v_i) : V^n \rightarrow V$ is uniquely defined by the following relation:

$$[\nabla_{i=0}^n(v_i)] v_m^n = v_m, \text{ for every } m \leq n.$$

Observe that, for every $m \leq n \leq t$, we have

$$f_t v_m^t = f_t w_n^t v_m^n = f_n v_m^n$$

so that, by the universal property of the direct sum, there exists a unique morphism $f : V \rightarrow X$ such that

$$(7) \quad f v_m = f_n v_m^n,$$

for any $m \in \mathbb{N}$, where $n \in \mathbb{N}$ and $m \leq n$. Thus

$$f_n v_m^n = f v_m = f [\nabla_{i=0}^n(v_i)] v_m^n \text{ for every } m \leq n.$$

By the universal property of $V^n := \oplus_{i=0}^n V_i$, f_n is the unique morphism that composed with v_m^n gives $f_n v_m^n$ for any $m \leq n$. We get that

$$f_n = f [\nabla_{i=0}^n(v_i)], \text{ for every } n \in \mathbb{N}.$$

In order to conclude that $V = \varinjlim V^i$, it remains to prove that $f : V \rightarrow X$ is the unique morphism with this property. Let $g : V \rightarrow X$ be a morphism such that $f_n = g [\nabla_{i=0}^n(v_i)]$ for every $n \in \mathbb{N}$. Then

$$f v_m = f_n v_m^n = g [\nabla_{i=0}^n(v_i)] v_m^n = g v_m \text{ for every } m, n \in \mathbb{N}, m \leq n.$$

By uniqueness of f with respect to (7), we get $g = f$. $\square$

COROLLARY 2.6. *Let (C, Δ, ε) be a coalgebra in a cocomplete abelian monoidal category $\mathcal{M}$ and let (M, ρ_M^r, ρ_M^l) be a C -bicomodule. Then*

$$(T_C^c(M), (\sigma_n)_{n \in \mathbb{N}}) = \varinjlim C^n(M).$$

Proof. Just observe that $C^n(M) = \oplus_{m=0}^{n-1} M^{\square m}$ and $\sigma_n = \nabla_{m=0}^{n-1}(i_m)$. $\square$

2.7. Note that $M^{\square n}$ is a C -bicomodule via $\rho_n^l := \rho_M^l \square M^{\square n-1}$ and $\rho_n^r := M^{\square n-1} \square \rho_M^r$.

Our next aim is to define, for any $n \in \mathbb{N} \setminus \{0\}$, a Hochschild 2-cocycle

$$\zeta^n : M^{\square n} \rightarrow C^n(M) \otimes C^n(M).$$

Then we will apply Lemma 2.3 to obtain that, for any $n > 0$, $C^{n+1}(M) = C^n(M) \oplus M^{\square n}$ can be endowed with a coalgebra structure $(C^{n+1}(M), \Delta_{\zeta^n}, \varepsilon_{\zeta^n})$ in $\mathcal{M}$ such that the canonical inclusion $\sigma_n^{n+1} : C^n(M) \rightarrow C^{n+1}(M)$ is a Hochschild extension of $C^n(M)$ with cokernel $M^{\square n}$. Then, by Proposition 1.8, $T_C^c(M)$ will carry a natural coalgebra structure that makes it the direct limit of $((C^i(M))_{i \in \mathbb{N}}, (\sigma_i^j)_{i,j \in \mathbb{N}})$ as a direct system of coalgebras.

Let $\chi_M : M \square M \rightarrow M \otimes M$ be the canonical inclusion and define

$$(8) \quad \zeta^1 = 0 \quad \text{and} \quad \zeta^n = - \sum_{t=1}^{n-1} (i_t^n \otimes i_{n-t}^n) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}), \forall n > 1.$$

where we identify $C \square X$ and $X \square C$ with X , for any C -bicomodule X .

PROPOSITION 2.8. *Let (C, Δ, ε) be a coalgebra in a cocomplete abelian monoidal category $\mathcal{M}$ and let (M, ρ_M^r, ρ_M^l) be a C -bicomodule. Let*

$$\begin{aligned} \Delta(1) &= \Delta, & \text{and} & \quad \varepsilon(1) = \varepsilon, \\ \bar{\rho}_1^l &= \rho_M^l, & \text{and} & \quad \bar{\rho}_1^r = \rho_M^r, \end{aligned}$$

and for every $n \geq 2$ set

$$\Delta(n) = \Delta_{\zeta^{n-1}}, \quad \text{and} \quad \varepsilon(n) = \varepsilon_{\zeta^{n-1}},$$

as defined in (5), (6), and let

$$\bar{\rho}_n^l = (\sigma_1^n \otimes M^{\square n})(\rho_M^l \square M^{\square n-1}), \quad \text{and} \quad \bar{\rho}_n^r = (M^{\square n} \otimes \sigma_1^n)(M^{\square n-1} \square \rho_M^r).$$

Then, for any $n \geq 1$, we have

- a) $(C^n(M), \Delta(n), \varepsilon(n))$ is a coalgebra.
- b) $(M^{\square n}, \bar{\rho}_n^l, \bar{\rho}_n^r)$ is a $C^n(M)$ -bicomodule such that the morphism $\zeta^n : M^{\square n} \rightarrow C^n(M) \otimes C^n(M)$, given by (8), defines a Hochschild 2-cocycle.
- c) $\varepsilon(n) = \varepsilon_C \pi_n^1$.
- d) For every $1 \leq t \leq n-1$ we have

$$(9) \quad \Delta(n) i_t^n = (i_t^n \otimes \sigma_t^n) \bar{\rho}_t^r + (\sigma_t^n \otimes i_t^n) \bar{\rho}_t^l - (\sigma_t^n \otimes \sigma_t^n) \zeta^t.$$

e) $\Delta(n)$ fulfils the following relations:

$$\begin{aligned}\Delta(n)i_0^n &= (i_0^n \otimes i_0^n) \Delta \\ \Delta(n)i_1^n &= (i_1^n \otimes i_0^n) \rho_M^r + (i_0^n \otimes i_1^n) \rho_M^l \\ \text{and for } 2 &\leq t \leq n-1 \\ \Delta(n)i_t^n &= (i_t^n \otimes i_0^n) (M^{\square t-1} \square \rho_M^r) + (i_0^n \otimes i_t^n) (\rho_M^l \square M^{\square t-1}) + \\ &\quad + \sum_{r=1}^{t-1} (i_r^n \otimes i_{t-r}^n) (M^{\square r-1} \square \chi_M \square M^{\square t-1-r}).\end{aligned}$$

Proof. Set $C^n = C^n(M)$ for any $n \geq 1$. Recall that $M^{\square n}$ is a C -bicomodule via

$$\rho_n^l := \rho_M^l \square M^{\square n-1}, \quad \text{and} \quad \rho_n^r := M^{\square n-1} \square \rho_M^r.$$

Let us prove all the statements of the theorem by induction on $n \geq 1$.

If $n = 1$, then $C^1 = C$ is a coalgebra and $M^{\square 1} = M$ is a C^1 -bicomodule by hypothesis. Obviously $\zeta^1 = 0$ fulfils (12). We have $\varepsilon(1) = \varepsilon = \varepsilon_C \pi_1^1$ and, since $i_0^1 = \text{Id}_C$ and $i_1^1 = 0$, we get

$$\Delta(1)i_0^1 = \Delta = (i_0^1 \otimes i_0^1) \Delta, \quad \Delta(1)i_1^1 = 0 = (i_1^1 \otimes i_0^1) \rho_M^r + (i_0^1 \otimes i_1^1) \rho_M^l.$$

Let $n \geq 2$. Assume all the assertions hold true for any $1 \leq t < n$.

Thus $(C^{n-1}, \Delta(n-1), \varepsilon(n-1))$ is a coalgebra in $\mathcal{M}$, $(M^{\square n-1}, \bar{\rho}_{n-1}^l, \bar{\rho}_{n-1}^r)$ is a C^{n-1} -bicomodule and ζ^{n-1} is a 2-cocycle. By Lemma 2.3 applied to " C " = C^{n-1} and " M " = $M^{\square n-1}$, then $(C^n, \Delta(n), \varepsilon(n))$ is a coalgebra. Moreover $\sigma_{t-1}^n : C^{t-1} \rightarrow C^t$ is a Hochschild extension of C^{t-1} with cokernel $M^{\square t-1}$ for any $1 \leq t < n$.

Since $(M^{\square n}, \rho_n^l, \rho_n^r)$ is a C -bicomodule and $\sigma_1^n : C \rightarrow C^n$ is a coalgebra homomorphism (as a composition of coalgebra homomorphism), then $(M^{\square n}, \bar{\rho}_n^l, \bar{\rho}_n^r)$ is a C^n -bicomodule, where

$$\bar{\rho}_n^l = (\sigma_1^n \otimes M^{\square n})(\rho_M^l \square M^{\square n-1}), \quad \text{and} \quad \bar{\rho}_n^r = (M^{\square n} \otimes \sigma_1^n)(M^{\square n-1} \square \rho_M^r).$$

Recall that, by definition, we have:

$$\begin{aligned}\Delta(n) &= \Delta_{\zeta^{n-1}} \\ &= (\sigma_{n-1}^n \otimes \sigma_{n-1}^n) \Delta(n-1) \pi_n^{n-1} + \left[\begin{array}{c} (i_{n-1}^n \otimes \sigma_{n-1}^n) \bar{\rho}_{n-1}^r + (\sigma_{n-1}^n \otimes i_{n-1}^n) \bar{\rho}_{n-1}^l + \\ - (\sigma_{n-1}^n \otimes \sigma_{n-1}^n) \zeta^{n-1} \end{array} \right] p_n^{n-1},\end{aligned}$$

For any $0 \leq t \leq n-1$, we have that

$$\begin{aligned}&[(i_{n-1}^n \otimes \sigma_{n-1}^n) \bar{\rho}_{n-1}^r + (\sigma_{n-1}^n \otimes i_{n-1}^n) \bar{\rho}_{n-1}^l - (\sigma_{n-1}^n \otimes \sigma_{n-1}^n) \zeta^{n-1}] p_n^{n-1} i_t^n \\ &= [(i_{n-1}^n \otimes \sigma_{n-1}^n) \bar{\rho}_{n-1}^r + (\sigma_{n-1}^n \otimes i_{n-1}^n) \bar{\rho}_{n-1}^l - (\sigma_{n-1}^n \otimes \sigma_{n-1}^n) \zeta^{n-1}] \delta_{t,n-1}\end{aligned}$$

so that we obtain

$$(10) \quad \Delta(n)i_t^n = (\sigma_{n-1}^n \otimes \sigma_{n-1}^n) \Delta(n-1) \pi_n^{n-1} i_t^n + \left[\begin{array}{c} (i_{n-1}^n \otimes \sigma_{n-1}^n) \bar{\rho}_{n-1}^r + (\sigma_{n-1}^n \otimes i_{n-1}^n) \bar{\rho}_{n-1}^l - \\ - (\sigma_{n-1}^n \otimes \sigma_{n-1}^n) \zeta^{n-1} \end{array} \right] \delta_{t,n-1}.$$

For $t = 0 < n-1$, we have:

$$\Delta(n)i_0^n = (\sigma_{n-1}^n \otimes \sigma_{n-1}^n) \Delta(n-1) i_0^{n-1},$$

so that, inductively we get:

$$\Delta(n)i_0^n = (\sigma_{n-1}^n \otimes \sigma_{n-1}^n) (i_0^{n-1} \otimes i_0^{n-1}) \Delta_C = (i_0^n \otimes i_0^n) \Delta_C.$$

Let us prove that for every $t \in \mathbb{N}$, such that $0 < t \leq n-1$ we have (9).

Now, we apply (10). If $t = n-1$, we get

$$\Delta(n)i_t^n = (i_{n-1}^n \otimes \sigma_{n-1}^n) \bar{\rho}_{n-1}^r + (\sigma_{n-1}^n \otimes i_{n-1}^n) \bar{\rho}_{n-1}^l - (\sigma_{n-1}^n \otimes \sigma_{n-1}^n) \zeta^{n-1}.$$

If $0 < t < n-1$, we get

$$\begin{aligned}\Delta(n)i_t^n &= (\sigma_{n-1}^n \otimes \sigma_{n-1}^n) \Delta(n-1) i_t^{n-1} \\ &= (\sigma_{n-1}^n \otimes \sigma_{n-1}^n) [(i_t^{n-1} \otimes \sigma_t^{n-1}) \bar{\rho}_t^r + (\sigma_t^{n-1} \otimes i_t^{n-1}) \bar{\rho}_t^l - (\sigma_t^{n-1} \otimes \sigma_t^{n-1}) \zeta^t] \\ &= (i_t^n \otimes \sigma_t^n) \bar{\rho}_t^r + (\sigma_t^n \otimes i_t^n) \bar{\rho}_t^l - (\sigma_t^n \otimes \sigma_t^n) \zeta^t.\end{aligned}$$

Thus we have obtained (9). Note that, for $t = 1$, by definition of $\bar{p}_1^l, \bar{p}_1^r$ and since $\zeta^1 = 0$, one gets

$$\Delta(n)i_1^n = (i_1^n \otimes \sigma_1^n) \bar{p}_1^r + (\sigma_1^n \otimes i_1^n) \bar{p}_1^l = (i_1^n \otimes i_0^n) \rho_M^r + (i_0^n \otimes i_1^n) \rho_M^l.$$

For $2 \leq t \leq n-1$, by definition of $\bar{p}_n^l, \bar{p}_n^r$ and of ζ^t , from (9), we get:

$$\begin{aligned} \Delta(n)i_t^n &= (i_t^n \otimes \sigma_t^n) (M^{\square t} \otimes \sigma_1^t) (M^{\square t-1} \square \rho_M^r) + (\sigma_t^n \otimes i_t^n) (\sigma_1^t \otimes M^{\square t}) (\rho_M^l \square M^{\square t-1}) + \\ &+ \sum_{r=1}^{t-1} (\sigma_t^n \otimes \sigma_t^n) (i_r^t \otimes i_{t-r}^t) (M^{\square r-1} \square \chi_M \square M^{\square t-1-r}), \end{aligned}$$

so that we obtain:

$$\begin{aligned} \Delta(n)i_t^n &= (i_t^n \otimes i_0^n) (M^{\square t-1} \square \rho_M^r) + (i_0^n \otimes i_t^n) (\rho_M^l \square M^{\square t-1}) + \\ &+ \sum_{r=1}^{t-1} (i_r^n \otimes i_{t-r}^n) (M^{\square r-1} \square \chi_M \square M^{\square t-1-r}). \end{aligned}$$

Moreover, by definition, we have:

$$(11) \quad \varepsilon(n) = \varepsilon_{\zeta^{n-1}} = \varepsilon(n-1) \pi_n^{n-1} + l_1[\varepsilon(n-1) \otimes \varepsilon(n-1)] \zeta^{n-1} p_n^{n-1}.$$

Since $n \geq 2$ and $\varepsilon(n-1) = \varepsilon_C \pi_{n-1}^1$, by (11), we have

$$\varepsilon(n) = \varepsilon_C \pi_{n-1}^1 \pi_n^{n-1} + l_1(\varepsilon_C \pi_{n-1}^1 \otimes \varepsilon_C \pi_{n-1}^1) \zeta^{n-1} p_n^{n-1} = \varepsilon_C \pi_n^1 + l_1(\varepsilon_C \otimes \varepsilon_C) (\pi_{n-1}^1 \otimes \pi_{n-1}^1) \zeta^{n-1} p_n^{n-1}.$$

By definition of ζ^{n-1} , if $n = 2$ we have $\zeta^1 = 0$ so that $(\pi_{n-1}^1 \otimes \pi_{n-1}^1) \zeta^{n-1} = 0$. If $n \geq 3$, we have

$$(\pi_{n-1}^1 \otimes \pi_{n-1}^1) \zeta^{n-1} = \sum_{t=1}^{n-2} (\pi_{n-1}^1 \otimes \pi_{n-1}^1) (i_t^{n-1} \otimes i_{n-1-t}^{n-1}) (M^{\square t-1} \square \chi_M \square M^{\square n-2-t}) = 0.$$

We conclude that $(\pi_n^1 \otimes \pi_n^1) \zeta^{n-1} = 0$ for any $n \geq 2$ and hence $\varepsilon(n) = \varepsilon_C \pi_n^1$.

Recall that ζ^n is a 2-cocycle means that it verifies the following relation:

$$(12) \quad 0 = b^2(\zeta^n) = (\zeta^n \otimes C^n) \bar{p}_n^r - [C^n \otimes \Delta(n)] \zeta^n + [\Delta(n) \otimes C^n] \zeta^n - (C^n \otimes \zeta^n) \bar{p}_n^l.$$

Now, since $\sigma_1^n = i_0^n$ and $n \geq 2$, we have

$$\begin{aligned} -(\zeta^n \otimes C^n) \bar{p}_n^r &= \sum_{t=1}^{n-1} [(i_t^n \otimes i_{n-t}^n) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \otimes C^n] (M^{\square n} \otimes \sigma_1^n) (M^{\square n-1} \square \rho_M^r) \\ &= \sum_{t=1}^{n-1} (i_t^n \otimes i_{n-t}^n \otimes \sigma_1^n) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t} \otimes C) (M^{\square n-1} \square \rho_M^r) \\ &= \sum_{t=1}^{n-1} (i_t^n \otimes i_{n-t}^n \otimes i_0^n) (M^{\square t} \otimes M^{\square n-1-t} \square \rho_M^r) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \end{aligned}$$

and

$$\begin{aligned} &- [C^n \otimes \Delta(n)] \zeta^n \\ &= [C^n \otimes \Delta(n)] \sum_{t=1}^{n-1} (i_t^n \otimes i_{n-t}^n) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \\ &= \sum_{t=1}^{n-1} (i_t^n \otimes \Delta(n) i_{n-t}^n) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \\ &= \sum_{t=1}^{n-1} \{ i_t^n \otimes \left[(i_{n-t}^n \otimes i_0^n) (M^{\square n-t-1} \square \rho_M^r) + (i_0^n \otimes i_{n-t}^n) (\rho_M^l \square M^{\square n-t-1}) + \right. \\ &\quad \left. + \sum_{r=1}^{n-t-1} (i_r^n \otimes i_{n-t-r}^n) (M^{\square r-1} \square \chi_M \square M^{\square n-t-1-r}) \right] \} (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \\ &= \sum_{t=1}^{n-1} (i_t^n \otimes i_{n-t}^n \otimes i_0^n) (M^{\square t} \otimes M^{\square n-t-1} \square \rho_M^r) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) + \\ &+ \sum_{t=1}^{n-1} (i_t^n \otimes i_0^n \otimes i_{n-t}^n) (M^{\square t} \otimes \rho_M^l \square M^{\square n-t-1}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) + \\ &+ \sum_{t=1}^{n-1} \sum_{r=1}^{n-t-1} (i_t^n \otimes i_r^n \otimes i_{n-t-r}^n) (M^{\square t} \otimes M^{\square r-1} \square \chi_M \square M^{\square n-t-1-r}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \\ &= -(\zeta^n \otimes C^n) \bar{p}_n^r + \sum_{t=1}^{n-1} (i_t^n \otimes i_0^n \otimes i_{n-t}^n) (M^{\square t} \otimes \rho_M^l \square M^{\square n-t-1}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) + \\ &+ \sum_{t=1}^{n-1} \sum_{r=1}^{n-t-1} (i_t^n \otimes i_r^n \otimes i_{n-t-r}^n) (M^{\square t} \otimes M^{\square r-1} \square \chi_M \square M^{\square n-t-1-r}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \end{aligned}$$

Analogously one has

$$-(C^n \otimes \zeta^n) \bar{p}_n^l = \sum_{t=1}^{n-1} (i_0^n \otimes i_t^n \otimes i_{n-t}^n) (\rho_M^l \square M^{\square t-1} \square M^{\square n-t}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t})$$

and

$$\begin{aligned} & [\Delta(n) \otimes C^n] \zeta^n \\ = & -(C^n \otimes \zeta^n) \overline{\rho}_n^l + \sum_{t=1}^{n-1} (i_t^n \otimes i_0^n \otimes i_{n-t}^n) (M^{\square t-1} \square \rho_M^r \otimes M^{\square n-t}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) + \\ & + \sum_{t=1}^{n-1} \sum_{r=1}^{t-1} (i_r^n \otimes i_{t-r}^n \otimes i_{n-t}^n) (M^{\square r-1} \square \chi_M \square M^{\square t-1-r} \otimes M^{\square n-t}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}). \end{aligned}$$

Now, for any $1 \leq t \leq n-1$, by definition of χ_M we have

$$\begin{aligned} & (i_t^n \otimes i_0^n \otimes i_{n-t}^n) (M^{\square t-1} \square \rho_M^r \otimes M^{\square n-t}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \\ = & (i_t^n \otimes i_0^n \otimes i_{n-t}^n) \left([M^{\square t-1} \square (\rho_M^r \otimes M) \chi_M \square M^{\square n-1-t}] \right) \\ = & (i_t^n \otimes i_0^n \otimes i_{n-t}^n) \left([M^{\square t-1} \square (M \otimes \rho_M^l) \chi_M \square M^{\square n-1-t}] \right) \\ = & (i_t^n \otimes i_0^n \otimes i_{n-t}^n) (M^{\square t} \otimes \rho_M^l \square M^{\square n-1-t}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \end{aligned}$$

an also

$$\begin{aligned} & \sum_{t=1}^{n-1} \sum_{r=1}^{t-1} (i_r^n \otimes i_{t-r}^n \otimes i_{n-t}^n) (M^{\square r-1} \square \chi_M \square M^{\square t-1-r} \otimes M^{\square n-t}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \\ = & \sum_{t=1}^{n-1} \sum_{r=1}^{t-1} (i_r^n \otimes i_{t-r}^n \otimes i_{n-t}^n) (M^{\square r} \otimes M^{\square t-1-r} \square \chi_M \square M^{\square n-1-t}) (M^{\square r-1} \square \chi_M \square M^{\square n-1-r}) \\ = & \sum_{t=1}^{n-1} \sum_{\substack{r+j=t \\ r,j>0}} (i_r^n \otimes i_j^n \otimes i_{n-t}^n) (M^{\square r} \otimes M^{\square j-1} \square \chi_M \square M^{\square n-1-t}) (M^{\square r-1} \square \chi_M \square M^{\square n-1-r}) \\ = & \sum_{\substack{r+j+k=n \\ k,r,j>0}} (i_r^n \otimes i_j^n \otimes i_k^n) (M^{\square r} \otimes M^{\square j-1} \square \chi_M \square M^{\square k-1}) (M^{\square r-1} \square \chi_M \square M^{\square n-1-r}) \\ = & \sum_{t=1}^{n-1} \sum_{\substack{r+k=t \\ r,k>0}} (i_t^n \otimes i_r^n \otimes i_k^n) (M^{\square t} \otimes M^{\square r-1} \square \chi_M \square M^{\square k-1}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \\ = & \sum_{t=1}^{n-1} \sum_{r=1}^{n-t-1} (i_t^n \otimes i_r^n \otimes i_{n-t-r}^n) (M^{\square t} \otimes M^{\square r-1} \square \chi_M \square M^{\square n-t-1-r}) (M^{\square t-1} \square \chi_M \square M^{\square n-1-t}) \end{aligned}$$

Then ζ^n satisfies (12). $\square$

THEOREM 2.9. *Let (C, Δ, ε) be a coalgebra in a cocomplete abelian monoidal category $\mathcal{M}$ with left exact tensor functors and let (M, ρ_M^r, ρ_M^l) be a C -bicomodule. $(T_C^c(M), (\sigma_i)_{i \in \mathbb{N}})$ carries a natural coalgebra structure that makes it the direct limit of $((C^i(M))_{i \in \mathbb{N}}, (\sigma_i^j)_{i,j \in \mathbb{N}})$ as a direct system of coalgebras.*

Proof. By Proposition 2.8, for any $n \in \mathbb{N} \setminus \{0\}$, the canonical inclusion $\sigma_{n-1}^n : C^{n-1}(M) \rightarrow C^n(M)$ is a Hochschild extension of $C^{n-1}(M)$ with cokernel $M^{\square n-1}$. In particular σ_{n-1}^n is a coalgebra homomorphism for any $n \in \mathbb{N}$. Then, by 2.4, σ_m^n is a coalgebra homomorphism for any $m, n \in \mathbb{N}$. Now, in view of Corollary 2.6 and Proposition 1.8, $T_C^c(M)$ carries a natural coalgebra structure that makes it the direct limit of $((C^i(M))_{i \in \mathbb{N}}, (\sigma_i^j)_{i,j \in \mathbb{N}})$ as a direct system of coalgebras. $\square$

LEMMA 2.10. *Let (C, Δ, ε) be a coalgebra in a cocomplete abelian monoidal category $\mathcal{M}$ and let (M, ρ_M^r, ρ_M^l) be a C -bicomodule. Let $T := T_C^c(M)$. Then, for any $m, n \geq 1$ the following relations hold true:*

$$\begin{aligned} (13) \quad & (p_m \otimes p_n) \Delta_T = (M^{\square m-1} \square \chi_M \square M^{\square n-1}) p_{m+n}, \text{ for any } m, n \geq 1; \\ (14) \quad & (p_m \otimes p_0) \Delta_T = (M^{\square m-1} \square \rho_M^r) p_m, \text{ for any } m \geq 1; \\ (15) \quad & (p_0 \otimes p_n) \Delta_T = (\rho_M^l \square M^{\square n-1}) p_n, \text{ for any } n \geq 1; \\ (16) \quad & (p_0 \otimes p_0) \Delta_T = \Delta_C p_0. \end{aligned}$$

Proof. By construction, the comultiplication Δ_T of T is uniquely defined by the following relation

$$\Delta_T \sigma_i = (\sigma_i \otimes \sigma_i) \Delta(i), \text{ for any } i \in \mathbb{N},$$

so that, for any $m, n, k \geq 0$ we have

$$\begin{aligned} (p_m \otimes p_n) \Delta_T i_k &= (p_m \otimes p_n) \Delta_T \sigma_{k+1} i_k^{k+1} \\ &= (p_m \otimes p_n) (\sigma_{k+1} \otimes \sigma_{k+1}) \Delta(k+1) i_k^{k+1} = (p_{k+1}^m \otimes p_{k+1}^n) \Delta(k+1) i_k^{k+1}. \end{aligned}$$

Recall that:

$$\begin{aligned} \Delta(k+1) i_0^{k+1} &= (i_0^{k+1} \otimes i_0^{k+1}) \Delta_C \\ \Delta(k+1) i_1^{k+1} &= (i_1^{k+1} \otimes i_0^{k+1}) \rho_M^r + (i_0^{k+1} \otimes i_1^{k+1}) \rho_M^l \\ \text{and for } 2 \leq t & \\ \Delta(k+1) i_t^{k+1} &= (i_t^{k+1} \otimes i_0^{k+1}) (M^{\square t-1} \square \rho_M^r) + (i_0^{k+1} \otimes i_t^{k+1}) (\rho_M^l \square M^{\square t-1}) + \\ &\quad + \sum_{r=1}^{t-1} (i_r^{k+1} \otimes i_{t-r}^{k+1}) (M^{\square r-1} \square \chi_M \square M^{\square t-1-r}). \end{aligned}$$

Then, for $k = 0, 1$ we have respectively

$$\begin{aligned} (p_m \otimes p_n) \Delta_T i_0 &= (p_1^m \otimes p_1^n) \Delta(1) i_0^1 = \begin{cases} 0, & \text{for } m \geq 1 \text{ or } n \geq 1; \\ \Delta(1) i_0^1, & \text{for } m = n = 0; \end{cases} \\ (p_m \otimes p_n) \Delta_T i_1 &= (p_2^m \otimes p_2^n) \Delta(2) i_1^2 \\ &= (p_2^m \otimes p_2^n) [(i_1^2 \otimes i_0^2) \rho_M^r + (i_0^2 \otimes i_1^2) \rho_M^l] = \begin{cases} 0, & \text{for } m \geq 1 \text{ and } n \geq 1; \\ \delta_{m,1} \rho_M^r, & \text{for } n = 0; \\ \delta_{n,1} \rho_M^l, & \text{for } m = 0; \end{cases} \end{aligned}$$

while, for $k \geq 2$, we have:

$$\begin{aligned} &(p_m \otimes p_n) \Delta_T i_k \\ &= (p_{k+1}^m \otimes p_{k+1}^n) \Delta(k+1) i_k^{k+1} \\ &= (p_{k+1}^m \otimes p_{k+1}^n) \left[(i_k^{k+1} \otimes i_0^{k+1}) (M^{\square k-1} \square \rho_M^r) + (i_0^{k+1} \otimes i_k^{k+1}) (\rho_M^l \square M^{\square k-1}) \right. \\ &\quad \left. + \sum_{r=1}^{k-1} (i_r^{k+1} \otimes i_{k-r}^{k+1}) (M^{\square r-1} \square \chi_M \square M^{\square k-1-r}) \right] \\ &= \begin{cases} \delta_{n,k-m} (M^{\square m-1} \square \chi_M \square M^{\square n-1}), & \text{for } m \geq 1 \text{ and } n \geq 1; \\ \delta_{0,k-m} (M^{\square k-1} \square \rho_M^r), & \text{for } n = 0; \\ \delta_{n,k} (\rho_M^l \square M^{\square k-1}), & \text{for } m = 0; \end{cases} \end{aligned}$$

so that, for any $k \geq 0$, we get:

$$\begin{aligned} (p_m \otimes p_n) \Delta_T i_k &= \delta_{n+m,k} (M^{\square m-1} \square \chi_M \square M^{\square n-1}) = (M^{\square m-1} \square \chi_M \square M^{\square n-1}) p_{m+n} i_k, \text{ for any } m, n \geq 1; \\ (p_m \otimes p_0) \Delta_T i_k &= \delta_{m,k} (M^{\square k-1} \square \rho_M^r) = (M^{\square m-1} \square \rho_M^r) p_m i_k, \text{ for any } m \geq 1; \\ (p_0 \otimes p_n) \Delta_T i_k &= \delta_{n,k} (\rho_M^l \square M^{\square k-1}) = (\rho_M^l \square M^{\square n-1}) p_n i_k, \text{ for any } n \geq 1; \\ (p_0 \otimes p_0) \Delta_T i_k &= \delta_{0,k} \Delta(1) = \Delta_C p_0 i_k. \end{aligned}$$

Therefore, we conclude. $\square$

PROPOSITION 2.11. *Let (C, Δ, ε) be a coalgebra in a cocomplete abelian monoidal category $\mathcal{M}$ and let (M, ρ_M^r, ρ_M^l) be a C -bicomodule. Let $T := T_C^c(M)$. Let E be a coalgebra and let $\alpha : E \rightarrow T$ and $\beta : E \rightarrow T$ be coalgebra homomorphisms. If $p_1 \alpha = p_1 \beta$, then $p_n \alpha = p_n \beta$ for any $n \geq 1$.*

Proof. Let us prove it by induction on $n \geq 1$, the case $n = 1$ being true by assumption. Thus let $n \geq 2$ be such that $p_n \alpha = p_n \beta$. Then

$$\begin{aligned} (\chi_M \square M^{\square n-1}) p_{n+1} \alpha &\stackrel{(13)}{=} (p_1 \otimes p_n) \Delta_T \alpha \\ &= (p_1 \alpha \otimes p_n \alpha) \Delta_T \\ &= (p_1 \beta \otimes p_n \beta) \Delta_T \\ &= (p_1 \otimes p_n) \Delta_T \beta \stackrel{(13)}{=} (\chi_M \square M^{\square n-1}) p_{n+1} \beta. \end{aligned}$$

Since χ_M is a monomorphism, then, by left exactness of the tensor functors, $\chi_M \square M^{\square n-1}$ is a monomorphism too, so that $p_{n+1} \alpha = p_{n+1} \beta$. $\square$

Our next aim is to prove the universal property of $T_C^c(M)$.

LEMMA 2.12. *Let (C, Δ, ε) be a coalgebra in a cocomplete abelian monoidal category $\mathcal{M}$ and let (M, ρ_M^r, ρ_M^l) be a C -bicomodule. Let $\delta : D \rightarrow E$ be a monomorphism which is a morphism of coalgebras such that the canonical morphism $\tilde{\delta} : \tilde{D} \rightarrow E$ of Notation 1.11 is a monomorphism. Then we have*

$$D^{\wedge_{\tilde{D}}}^n = (D^{\wedge_E^n}, \xi_n).$$

Proof. Denote by (L, p) the cokernel of $\delta : D \rightarrow E$ in $\mathcal{M}$ and by $(\tilde{L}, \tilde{p})$ the cokernel of $\xi_1 : D \rightarrow \tilde{D}$ in $\mathcal{M}$. Since $p\tilde{\delta}\xi_1 = p\delta = 0$, by the universal property of the cokernel, there exists a unique morphism $\lambda : \tilde{L} \rightarrow L$ such that

$$\lambda\tilde{p} = p\tilde{\delta}.$$

Since $\tilde{\delta}$ is a monomorphism, so is λ . Therefore, we have that

$$D^{\wedge_{\tilde{D}}}^n := \text{Ker}(\tilde{p}^{\otimes n} \Delta_{\tilde{D}}^{n-1}) = \text{Ker}(\lambda^{\otimes n} \tilde{p}^{\otimes n} \Delta_{\tilde{D}}^{n-1}) = \text{Ker}(p^{\otimes n} \tilde{\delta}^{\otimes n} \Delta_{\tilde{D}}^{n-1}) = \text{Ker}(p^{\otimes n} \Delta_E^{n-1} \tilde{\delta}).$$

We conclude by observing that, since, by definition, $(D^{\wedge_E^n}, \delta_n = \tilde{\delta}\xi_n) := \text{Ker}(p^{\otimes n} \Delta_E^{n-1})$ where $\tilde{\delta}$ is a monomorphism, the following relation holds true

$$(D^{\wedge_E^n}, \xi_n) = \text{Ker}(p^{\otimes n} \Delta_E^{n-1} \tilde{\delta}).$$

□

THEOREM 2.13. *Let (C, Δ, ε) be a coalgebra in a cocomplete abelian monoidal category $\mathcal{M}$ and let (M, ρ_M^r, ρ_M^l) be a C -bicomodule. Let $\delta : D \rightarrow E$ be a monomorphism which is a homomorphism of coalgebras such that the canonical morphism $\tilde{\delta} : \tilde{D} \rightarrow E$ of Notation 1.11 is a monomorphism. Let $f_C : \tilde{D} \rightarrow C$ be a coalgebra homomorphism and let $f_M : \tilde{D} \rightarrow M$ be a morphism of C -bicomodules such that $f_M \xi_1 = 0$, where $\tilde{D}$ is a C -bicomodule via f_C . Then there is a unique morphism $f : \tilde{D} \rightarrow T_C^c(M)$ such that*

$$f\xi_n = \sigma_n f_n, \text{ for any } n \in \mathbb{N},$$

where

$$(17) \quad f_n = \sum_{t=0}^n i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \xi_n$$

and $\overline{\Delta}_{\tilde{D}}^n : \tilde{D} \rightarrow \tilde{D}^{\square n+1}$ is the n^{th} -iteration of $\overline{\Delta}_{\tilde{D}}$ ($\overline{\Delta}_{\tilde{D}}^{-1} = f_C, \overline{\Delta}_{\tilde{D}}^0 = \text{Id}_{\tilde{D}}, \overline{\Delta}_{\tilde{D}}^1 = \overline{\Delta}_{\tilde{D}} : \tilde{D} \rightarrow \tilde{D} \square \tilde{D}$). Moreover:

1) f is a coalgebra homomorphism;

2) $p_0 f = f_C$ and $p_1 f = f_M$, where $p_n : T_C^c(M) \rightarrow M^{\square n}$ denotes the canonical projection.

Furthermore, any coalgebra homomorphism $f : \tilde{D} \rightarrow T_C^c(M)$ that fulfils 2) satisfies the following relation:

$$(18) \quad p_k f = f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1} \text{ for any } k \in \mathbb{N}.$$

Proof. Set $T := T_C^c(M)$. Following (1.9), denote by (L, p) the cokernel of $\delta : D \rightarrow E$ in $\mathcal{M}$ and let

$$(D^{\wedge_E^n}, \delta_n) := \text{Ker}(p^{\otimes n} \Delta_E^{n-1})$$

for any $n \in \mathbb{N} \setminus \{0\}$, where $\Delta^n : E \rightarrow E^{\otimes n+1}$ is the n^{th} iterated comultiplication of E ($\Delta^0 = \text{Id}_E, \Delta^1 := \Delta_E$). Since $f_C \xi_1 : D \rightarrow C$ is a coalgebra homomorphism, then D becomes a C -bicomodule and ξ_1 a morphism of C -bicomodules. Set $C^n = C^n(M)$ and $D^n = D^{\wedge_E^n}$ for any $n \in \mathbb{N} \setminus \{0\}$. Define $f_n : D^n \rightarrow C^n$, for every $n \in \mathbb{N}$, as in (17).

Note that, for every $n \geq 1$, $i_n^n = 0$ so that we have

$$f_n = \sum_{t=0}^{n-1} i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \xi_n.$$

Let us prove that $f_{n+1}\xi_n^{n+1} = \sigma_n^{n+1}f_n$ for any $n \in \mathbb{N}$. We have that

$$\begin{aligned} f_{n+1}\xi_n^{n+1} &= \sum_{t=0}^n i_t^{n+1} f_M^{\square t} \overline{\Delta_D}^{t-1} \xi_{n+1} \xi_n^{n+1} \\ &= \sum_{t=0}^n i_t^{n+1} f_M^{\square t} \overline{\Delta_D}^{t-1} \xi_n \\ &= \sum_{t=0}^{n-1} \sigma_n^{n+1} i_t^n f_M^{\square t} \overline{\Delta_D}^{t-1} \xi_n + i_n^{n+1} f_M^{\square n} \overline{\Delta_D}^{n-1} \xi_n \\ &= \sigma_n^{n+1} f_n + i_n^{n+1} f_M^{\square n} \overline{\Delta_D}^{n-1} \xi_n. \end{aligned}$$

Let $(\tilde{L}, \tilde{p})$ be the cokernel of $\xi_1 : D \rightarrow \tilde{D}$ in $\mathcal{M}$. Let $\chi_{\tilde{L}} : \tilde{L} \square \tilde{L} \rightarrow \tilde{L} \otimes \tilde{L}$ be the canonical injection. Define $\chi_{\tilde{L}}^n : \tilde{L}^{\square n} \rightarrow \tilde{L}^{\otimes n}$, for every $n \in \mathbb{N}$, by setting $\chi_{\tilde{L}}^0 = \text{Id}_C$, $\chi_{\tilde{L}}^1 = \text{Id}_{\tilde{L}}$, $\chi_{\tilde{L}}^2 = \chi_{\tilde{L}}$ and $\chi_{\tilde{L}}^n = (\tilde{L}^{\otimes n-2} \otimes \chi_{\tilde{L}})(\chi_{\tilde{L}}^{n-1} \square \tilde{L})$ for any $n > 2$. Since the tensor functors are left exact, $\chi_{\tilde{L}}^n$ is a monomorphism. By Lemma 2.12, we have

$$(D^{\wedge_E^n}, \xi_n) = D^{\wedge_{\tilde{D}}^n} = \text{Ker}(\tilde{p}^{\otimes n} \Delta_{\tilde{D}}^{n-1})$$

Thus, for any $n \geq 1$, we have

$$\text{Ker}(\tilde{p}^{\otimes n} \Delta_{\tilde{D}}^{n-1}) = \text{Ker}(\chi_{\tilde{L}}^n \circ \tilde{p}^{\square n} \overline{\Delta_D}^{n-1}) = \text{Ker}(\tilde{p}^{\square n} \overline{\Delta_D}^{n-1}).$$

so that we get

$$(19) \quad \tilde{p}^{\square n} \overline{\Delta_D}^{n-1} \xi_n = 0.$$

Now, since $f_M \xi_1 = 0$, there is a unique morphism of C -bicomodules $\lambda : \tilde{L} \rightarrow M$ such that $\lambda \tilde{p} = f_M$. Thus:

$$(20) \quad f_M^{\square n} \overline{\Delta_D}^{n-1} \xi_k = \lambda^{\square n} (\tilde{p}^{\square n} \overline{\Delta_D}^{n-1} \xi_n) \xi_k \stackrel{(19)}{=} 0, \text{ for any } k \leq n.$$

We conclude that $f_{n+1}\xi_n^{n+1} = \sigma_n^{n+1}f_n$ for any $n \in \mathbb{N}$ so that $(\sigma_n f_n : D^n \rightarrow T)_n$ is a compatible family of morphisms in $\mathcal{M}$. Thus there is a unique morphism $f : \tilde{D} \rightarrow T$ such that

$$f \xi_k = \sigma_k f_k, \text{ for any } k \in \mathbb{N}.$$

We have that

$$(21) \quad p_k^n \sum_{t=0}^k i_t^k f_M^{\square t} \overline{\Delta_D}^{t-1} \xi_k = f_M^{\square n} \overline{\Delta_D}^{n-1} \xi_k$$

for any $0 \leq n < k$. Note that, for $k \leq n$, $p_k^n = 0$ and the right side of (21) is zero by (20). Thus, the relation above holds true for any $k, n \in \mathbb{N}$. Then we get:

$$p_n f \xi_k = p_n \sigma_k f_k = p_k^n f_k = p_k^n \sum_{t=0}^k i_t^k f_M^{\square t} \overline{\Delta_D}^{t-1} \xi_k = f_M^{\square n} \overline{\Delta_D}^{n-1} \xi_k, \text{ for any } k, n \in \mathbb{N}.$$

We conclude that $p_n f = f_M^{\square n} \overline{\Delta_D}^{n-1}$, for any $n \in \mathbb{N}$. In particular, for $n = 0, 1$, we get $p_0 f = \overline{\Delta_D}^{-1} = f_C$ and $p_1 f = f_M \overline{\Delta_D}^0 = f_M$.

We have now to prove that f is a coalgebra homomorphism. Let us check that $f_n : D^n \rightarrow C^n$ is a coalgebra homomorphism for every $n \in \mathbb{N}$.

For $n = 0$, $f_n = 0$ and there is nothing to prove.

Assume $n \geq 1$. By Proposition 2.8, we get:

$$\begin{aligned}
& \Delta(n)f_n \\
&= \sum_{t=0}^n \Delta(n)i_t^n f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= \Delta(n)i_0^n f_M^{\square 0} \bar{\Delta}_{\tilde{D}}^{-1} \xi_n + \Delta(n)i_1^n f_M^{\square 1} \bar{\Delta}_{\tilde{D}}^0 \xi_n + \sum_{t=2}^n \Delta(n)i_t^n f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= \Delta(n)i_0^n f_C \xi_n + \Delta(n)i_1^n f_M \xi_n + \sum_{t=2}^n \Delta(n)i_t^n f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= (i_0^n \otimes i_0^n) \Delta_C f_C \xi_n + [(i_1^n \otimes i_0^n) \rho_M^r + (i_0^n \otimes i_1^n) \rho_M^l] f_M \xi_n + \\
&\quad + \sum_{t=2}^n \left[\frac{(i_t^n \otimes i_0^n) (M^{\square t-1} \square \rho_M^r) + (i_0^n \otimes i_t^n) (\rho_M^l \square M^{\square t-1})}{\sum_{r=1}^{t-1} (i_r^n \otimes i_{t-r}^n) (M^{\square r-1} \square \chi_M \square M^{\square t-1-r})} \right] f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= (i_0^n f_C \otimes i_0^n f_C) \Delta_{\tilde{D}} \xi_n + [(i_1^n \otimes i_0^n) \rho_M^r + (i_0^n \otimes i_1^n) \rho_M^l] f_M \xi_n + \\
&\quad + \sum_{t=2}^n \left[(i_t^n \otimes i_0^n) (M^{\square t-1} \square \rho_M^r) + (i_0^n \otimes i_t^n) (\rho_M^l \square M^{\square t-1}) \right] f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n + \\
&\quad + \sum_{t=2}^n \sum_{r=1}^{t-1} (i_r^n \otimes i_{t-r}^n) (M^{\square r-1} \square \chi_M \square M^{\square t-1-r}) f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= (i_0^n f_C \otimes i_0^n f_C) \Delta_{\tilde{D}} \xi_n + \\
&\quad + \sum_{t=1}^n \left[(i_t^n \otimes i_0^n) (M^{\square t-1} \square \rho_M^r) + (i_0^n \otimes i_t^n) (\rho_M^l \square M^{\square t-1}) \right] f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n + \\
&\quad + \sum_{t=2}^n \sum_{r=1}^{t-1} (i_r^n \otimes i_{t-r}^n) (M^{\square r-1} \square \chi_M \square M^{\square t-1-r}) f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n.
\end{aligned}$$

Now:

$$\begin{aligned}
\sum_{t=1}^n (i_t^n \otimes i_0^n) (M^{\square t-1} \square \rho_M^r) f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n &= \sum_{t=1}^n (i_t^n \otimes i_0^n) (f_M^{\square t-1} \square \rho_M^r f_M) \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= \sum_{t=1}^n (i_t^n \otimes i_0^n) [f_M^{\square t-1} \square (f_M \otimes C) \rho_M^r] \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= \sum_{t=1}^n (i_t^n \otimes i_0^n) (f_M^{\square t-1} \square f_M \otimes C) (\tilde{D}^{\square t-1} \square \rho_{\tilde{D}}^r) \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= \sum_{t=1}^n (i_t^n \otimes i_0^n) (f_M^{\square t} \otimes C) [\tilde{D}^{\square t-1} \square (\tilde{D} \otimes f_C) \Delta_{\tilde{D}}] \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= \sum_{t=1}^n (i_t^n \otimes i_0^n) (f_M^{\square t} \otimes f_C) [\tilde{D}^{\square t-1} \square \Delta_{\tilde{D}}] \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= \sum_{t=1}^n (i_t^n \otimes i_0^n) (f_M^{\square t} \otimes f_C) [\bar{\Delta}_{\tilde{D}}^{t-1} \otimes \tilde{D}] \Delta_{\tilde{D}} \xi_n \\
&= \sum_{t=1}^n (i_t^n f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \otimes i_0^n f_C) \Delta_{\tilde{D}} \xi_n
\end{aligned}$$

Analogously one gets

$$\sum_{t=1}^n (i_0^n \otimes i_t^n) (\rho_M^l \square M^{\square t-1}) f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n = \sum_{t=1}^n (i_0^n f_C \otimes i_t^n f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1}) \Delta_{\tilde{D}} \xi_n$$

Moreover we have:

$$\begin{aligned}
& \sum_{t=2}^n \sum_{r=1}^{t-1} (i_r^n \otimes i_{t-r}^n) (M^{\square r-1} \square \chi_M \square M^{\square t-1-r}) f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= \sum_{t=2}^n \sum_{r=1}^{t-1} (i_r^n \otimes i_{t-r}^n) (f_M^{\square r} \otimes f_M^{\square t-r}) (\tilde{D}^{\square r-1} \square \chi_{\tilde{D}} \square \tilde{D}^{\square t-1-r}) \bar{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= \sum_{t=2}^n \sum_{r=1}^{t-1} (i_r^n f_M^{\square r} \otimes i_{t-r}^n f_M^{\square t-r}) (\bar{\Delta}_{\tilde{D}}^{r-1} \otimes \bar{\Delta}_{\tilde{D}}^{t-1-r}) \Delta_{\tilde{D}} \xi_n \\
&= \sum_{t=2}^n \sum_{r=1}^{t-1} (i_r^n f_M^{\square r} \bar{\Delta}_{\tilde{D}}^{r-1} \otimes i_{t-r}^n f_M^{\square t-r} \bar{\Delta}_{\tilde{D}}^{t-1-r}) \Delta_{\tilde{D}} \xi_n
\end{aligned}$$

So we get

$$\begin{aligned}
\Delta(n)f_n &= (i_0^n f_C \otimes i_0^n f_C) \Delta_{\tilde{D}} \xi_n + \\
&\quad + \sum_{t=1}^n (i_t^n f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1} \otimes i_0^n f_C) \Delta_{\tilde{D}} \xi_n + \sum_{t=1}^n (i_0^n f_C \otimes i_t^n f_M^{\square t} \bar{\Delta}_{\tilde{D}}^{t-1}) \Delta_{\tilde{D}} \xi_n + \\
&\quad + \sum_{t=2}^n \sum_{r=1}^{t-1} (i_r^n f_M^{\square r} \bar{\Delta}_{\tilde{D}}^{r-1} \otimes i_{t-r}^n f_M^{\square t-r} \bar{\Delta}_{\tilde{D}}^{t-1-r}) \Delta_{\tilde{D}} \xi_n
\end{aligned}$$

On the other hand we have

$$\begin{aligned}
& (f_n \otimes f_n) \Delta_{D^n} \\
&= \left(\sum_{t=0}^n i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \xi_n \otimes \sum_{k=0}^n i_k^n f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1} \xi_n \right) \Delta_{D^n} \\
&= \left(\sum_{t=0}^n i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \otimes \sum_{k=0}^n i_k^n f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1} \right) \Delta_{\tilde{D}} \xi_n \\
&= (i_0^n f_M^{\square 0} \overline{\Delta}_{\tilde{D}}^{-1} \otimes i_0^n f_M^{\square 0} \overline{\Delta}_{\tilde{D}}^{-1}) \Delta_{\tilde{D}} \xi_n + \\
&\quad + \sum_{t=1}^n (i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \otimes i_0^n f_M^{\square 0} \overline{\Delta}_{\tilde{D}}^{-1}) \Delta_{\tilde{D}} \xi_n + \sum_{k=1}^n (i_0^n f_M^{\square 0} \overline{\Delta}_{\tilde{D}}^{-1} \otimes i_k^n f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1}) \Delta_{\tilde{D}} \xi_n \\
&\quad + \left(\sum_{t=1}^n i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \otimes \sum_{k=1}^n i_k^n f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1} \right) \Delta_{\tilde{D}} \xi_n \\
&= (i_0^n f_C \otimes i_0^n f_C) \Delta_{\tilde{D}} \xi_n + \sum_{t=1}^n (i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \otimes i_0^n f_C) \Delta_{\tilde{D}} \xi_n + \sum_{k=1}^n (i_0^n f_C \otimes i_k^n f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1}) \Delta_{\tilde{D}} \xi_n + \\
&\quad + \sum_{t=1}^n \sum_{k=1}^n (i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \otimes i_k^n f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1}) \Delta_{\tilde{D}} \xi_n \\
&= \Delta(n) f_n - \sum_{t=2}^n \sum_{r=1}^{t-1} (i_r^n f_M^{\square r} \overline{\Delta}_{\tilde{D}}^{r-1} \otimes i_{t-r}^n f_M^{\square t-r} \overline{\Delta}_{\tilde{D}}^{t-1-r}) \Delta_{\tilde{D}} \xi_n + \\
&\quad + \sum_{t=1}^n \sum_{k=1}^n (i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \otimes i_k^n f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1}) \Delta_{\tilde{D}} \xi_n.
\end{aligned}$$

But

$$\begin{aligned}
& (i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \otimes i_k^n f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1}) \Delta_{\tilde{D}} \xi_n \\
&= (i_t^n f_M^{\square t} \otimes i_k^n f_M^{\square k}) (\overline{\Delta}_{\tilde{D}}^{t-1} \otimes \overline{\Delta}_{\tilde{D}}^{k-1}) \Delta_{\tilde{D}} \xi_n \\
&= (i_t^n f_M^{\square t} \otimes i_k^n f_M^{\square k}) (\tilde{D}^{\square t-1} \square \chi_{\tilde{D}} \square \tilde{D}^{\square k-1}) \overline{\Delta}_{\tilde{D}}^{t+k-1} \xi_n \\
&= (i_t^n \otimes i_k^n) (f_M^{\square t} \otimes f_M^{\square k}) (\tilde{D}^{\square t-1} \square \chi_{\tilde{D}} \square \tilde{D}^{\square k-1}) \overline{\Delta}_{\tilde{D}}^{t+k-1} \xi_n \\
&= (i_t^n \otimes i_k^n) (M^{\square t-1} \square \chi_M \square M^{\square k-1}) f_M^{\square t+k} \overline{\Delta}_{\tilde{D}}^{t+k-1} \xi_n.
\end{aligned}$$

By (20), the last term is zero whenever $t+k > n$, so that:

$$\begin{aligned}
\sum_{t=1}^n \sum_{k=1}^n (i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \otimes i_k^n f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1}) \Delta_{\tilde{D}} \xi_n &= \sum_{\substack{1 \leq t, k \leq n \\ t+k \leq n}} (i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \otimes i_k^n f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1}) \Delta_{\tilde{D}} \xi_n \\
&= \sum_{t=2}^n \sum_{r=1}^{t-1} (i_r^n f_M^{\square r} \overline{\Delta}_{\tilde{D}}^{r-1} \otimes i_{t-r}^n f_M^{\square t-r} \overline{\Delta}_{\tilde{D}}^{t-1-r}) \Delta_{\tilde{D}} \xi_n,
\end{aligned}$$

and hence

$$(f_n \otimes f_n) \Delta_{D^n} = \Delta(n) f_n.$$

Furthermore $\varepsilon(0) f_0 = 0 = \varepsilon_{D^0}$, while, for every $n \geq 1$, we have

$$\begin{aligned}
\varepsilon(n) f_n &= \sum_{t=0}^n \varepsilon(n) i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \xi_n \\
&= \sum_{t=0}^n \varepsilon_C \pi_n^1 i_t^n f_M^{\square t} \overline{\Delta}_{\tilde{D}}^{t-1} \xi_n = \varepsilon_C \pi_n^1 i_0^n f_M^{\square 0} \overline{\Delta}_{\tilde{D}}^{0-1} \xi_n = \varepsilon_C i_0^1 f_C \xi_n = \varepsilon_C f_C \xi_n = \varepsilon_{D^n}.
\end{aligned}$$

We conclude that f_n is a coalgebra homomorphism. Now, by construction, f is the unique morphism such that $f \xi_k = \sigma_k f_k$, for any $k \in \mathbb{N}$. By Proposition 1.10, $((D^{\wedge^c})_{i \in \mathbb{N}}, (\xi_i^j)_{i, j \in \mathbb{N}})$ is a direct system in $\mathcal{M}$ whose direct limit $\tilde{D}$ carries a natural coalgebra structure that makes it the direct limit of $((D^{\wedge^c})_{i \in \mathbb{N}}, (\xi_i^j)_{i, j \in \mathbb{N}})$ as a direct system of coalgebras. Since σ_k is a coalgebra homomorphism so is $\sigma_k f_k$ and hence f is a coalgebra homomorphism.

Assume now that $g : E \rightarrow T$ is another coalgebra homomorphism such that $p_0 g = f_C$ and $p_1 g = f_M$. Then, by Proposition 2.11, we have $p_n g = p_n f$ for any $n \in \mathbb{N}$. $\square$

LEMMA 2.14. *Let $(X_i)_{i \in \mathbb{N}}$ be a family of objects in a cocomplete and complete abelian category $\mathcal{M}$ satisfying AB5. Let Y be an object in $\mathcal{M}$ and $f : Y \rightarrow \oplus_{i \in \mathbb{N}} X_i$ be a morphism such that*

$$p_k f = 0 \text{ for any } k \in \mathbb{N},$$

where $p_k : \oplus X_i \rightarrow X_k$ denotes the canonical projection. Then $f = 0$.

Proof. By [Po, Corollary 8.10, page 61], $\mathcal{M}$ is a C_2 -category so that the conclusion follows by [Po, Proposition, page 54]. $\square$

THEOREM 2.15. *Let (C, Δ, ε) be a coalgebra in a cocomplete and complete abelian monoidal category $\mathcal{M}$ satisfying AB5. Let (M, ρ_M^r, ρ_M^l) be a C -bicomodule. Let $\delta : D \rightarrow E$ be a monomorphism which is a homomorphism of coalgebras. Let $f_C : \tilde{D} \rightarrow C$ be a coalgebra homomorphism and let $f_M : \tilde{D} \rightarrow M$ be a morphism of C -bicomodules such that $f_M \xi_1 = 0$, where $\tilde{D}$ is a bicomodule via f_C . Then there is a unique coalgebra homomorphism $f : \tilde{D} \rightarrow T_C^c(M)$ such that $p_0 f = f_C$ and $p_1 f = f_M$, where $p_n : T_C^c(M) \rightarrow M^{\square n}$ denotes the canonical projection.*

$$\begin{array}{ccccc}
 T_C^c(M) & \xrightarrow{p_1} & M & & \\
 p_0 \downarrow & \nearrow f & \uparrow f_M & \searrow \theta & \\
 C & \xleftarrow{f_C} & \tilde{D} & \xleftarrow{\xi_1} & D
 \end{array}$$

Proof. Since $\mathcal{M}$ satisfies AB5, the morphism $\tilde{\delta} : \tilde{D} \rightarrow E$ of Notation 1.11 is a monomorphism, so that, by applying Theorem 2.13, there is a coalgebra homomorphism $f : \tilde{D} \rightarrow T_C^c(M)$ such that $p_0 f = f_C$ and $p_1 f = f_M$. Furthermore any such a coalgebra homomorphism fulfils the following relation

$$p_k f = f_M^{\square k} \overline{\Delta}_{\tilde{D}}^{k-1} \text{ for any } k \in \mathbb{N},$$

where $\overline{\Delta}_{\tilde{D}}^n : \tilde{D} \rightarrow \tilde{D}^{\square n+1}$ is the n^{th} -iteration of $\overline{\Delta}_{\tilde{D}}$ as defined in Theorem 2.13.

Now, let $f, g : \tilde{D} \rightarrow T$ be morphisms such that

$$p_k f = p_k g \text{ for any } k \in \mathbb{N}.$$

By Lemma 2.14 $f = g$. $\square$

LEMMA 2.16. *Let E be a coalgebra in an abelian monoidal category $\mathcal{M}$. Let $f : E \rightarrow L$ and $g : E \rightarrow M$ be morphism in $\mathcal{M}$. Then*

$$\text{Ker}(f) \wedge_E \text{Ker}(g) = \text{Ker}[(f \otimes g) \circ \triangle_E].$$

Proof. Let $(X, i_X) = \text{Ker}(f)$ and let $(Y, i_Y) = \text{Ker}(g)$. Let $p_X : E \rightarrow E/X$ and $p_Y : E \rightarrow E/Y$ be the canonical quotient maps. Since $f i_X = 0$, by the universal property of the cokernel, there exists a unique morphism $\gamma_X : E/X \rightarrow L$ such that $\gamma_X p_X = f$. Moreover, we have $(E/X, p_X) = \text{coker}(i_X) = \text{coker}(\text{Ker}(f)) = \text{coim}(f)$. As $\mathcal{M}$ is abelian, it is straightforward to prove that $(E/X, \gamma_X) = \text{Im}(f)$. In particular γ_X is a monomorphism. Analogously one gets a monomorphism $\gamma_Y : E/Y \rightarrow M$ such that $\gamma_Y p_Y = g$. Since $\mathcal{M}$ has left exact tensor functors, then $\gamma_X \otimes \gamma_Y$ is a monomorphism, so that, by definition, we get:

$$X \wedge_E Y := \text{Ker}[(p_X \otimes p_Y) \circ \triangle_E] = \text{Ker}[(\gamma_X \otimes \gamma_Y)(p_X \otimes p_Y) \circ \triangle_E] = \text{Ker}[(f \otimes g) \circ \triangle_E].$$

$\square$

PROPOSITION 2.17. *Let $\delta : D \rightarrow C$ be a monomorphism which is a coalgebra homomorphism in an abelian monoidal category $\mathcal{M}$. Then we have*

$$(22) \quad D^{\wedge_C^m} \wedge_C D^{\wedge_C^n} = D^{\wedge_C^{m+n}}.$$

Proof. Set $(L, p) = \text{coker}(\sigma)$. Let

$$(D^{\wedge_C^n}, \delta_n) := \text{Ker}(p^{\otimes n} \Delta_C^{n-1}).$$

By Lemma 2.16, we have:

$$D^{\wedge_C^{m+n}} = \text{Ker}[p^{\otimes m+n} \Delta_C^{m+n-1}] = \text{Ker}[(p^{\otimes m} \Delta_C^{m-1} \otimes p^{\otimes n} \Delta_C^{n-1}) \Delta_C] = D^{\wedge_C^m} \wedge_C D^{\wedge_C^n}.$$

$\square$

THEOREM 2.18. *Let (C, Δ, ε) be a coalgebra in a cocomplete and complete abelian monoidal category $\mathcal{M}$ satisfying AB5. Let (M, ρ_M^r, ρ_M^l) be a C -bicomodule. Let $T := T_C^c(M)$. Then*

$$(C^n(M), \sigma_n) = C^{\wedge_T^n},$$

for every $n \in \mathbb{N}$.

Proof. Let $n \in \mathbb{N}$. Let

$$\lambda_a^n : M^{\square a} \rightarrow \oplus_{b \geq n} M^{\square b}$$

be defined by

$$\lambda_a^n = \begin{cases} \text{the canonical injection} & \text{if } a \geq n, \\ 0 & \text{otherwise.} \end{cases}$$

Define

$$\nu_n : \oplus_{a \geq n} M^{\square a} \rightarrow T$$

as the codiagonal map of the family $(i_a)_{a \geq n}$ so that we have

$$\nu_n \circ \lambda_a^n = \begin{cases} i_a & \text{for every } a \geq n \\ 0 & \text{otherwise.} \end{cases}$$

Define

$$\tau_n : T \rightarrow \oplus_{a \geq n} M^{\square a}$$

as the codiagonal map of the family $(\lambda_a^n)_{a \in \mathbb{N}}$, that is

$$\tau_n i_a = \lambda_a^n, \text{ for every } a \in \mathbb{N}.$$

Thus, we have

$$\tau_n \nu_n \lambda_a^n = \tau_n i_a = \lambda_a^n \text{ for } a \geq n,$$

so that

$$\tau_n \nu_n = \text{Id}_{\oplus_{a \geq n} M^{\square a}} \text{ for every } n \in \mathbb{N}.$$

Let $C^n = C^n(M)$ for every $n \in \mathbb{N}$. Let us prove the following sequence

$$0 \rightarrow C^n \xrightarrow{\sigma_n} T \xrightarrow{\tau_n} \oplus_{a \geq n} M^{\square a} \rightarrow 0$$

is exact. We check that $(\oplus_{a \geq n} M^{\square a}, \tau_n) = \text{Coker}(\sigma_n)$.

Since $\tau_n \nu_n = \text{Id}$, it is clear that τ_n is an epimorphism and that ν_n is a monomorphism.

From

$$\tau_n \sigma_n i_a^n = \tau_n i_a = \lambda_a^n = 0, \text{ for every } 0 \leq a \leq n-1,$$

we deduce that

$$\tau_n \sigma_n = 0 \text{ for every } n \in \mathbb{N}.$$

Let $f : T \rightarrow X$ be a morphism such that $f \sigma_n = 0$ for every $n \in \mathbb{N}$. Thus, for every $0 \leq a \leq n-1$, we have

$$f i_a = f \sigma_n i_a^n = 0$$

Set

$$\bar{f} = f \nu_n$$

and let us prove that $f = \bar{f} \tau_n$. From

$$\bar{f} \tau_n i_a = f \nu_n \lambda_a^n = \begin{cases} f i_a & \text{for every } a \geq n \\ 0 & \text{otherwise.} \end{cases}$$

we deduce that $\bar{f} \tau_n i_a = f i_a$, for every $a \in \mathbb{N}$, and hence $\bar{f} \tau_n = f$.

Let us prove that $C^n = C^{\wedge_T^n}$, for every $n \in \mathbb{N}$.

The case $n = 0$ is trivial. Let us prove the equality above for every $n \geq 1$ by induction on n .

If $n = 1$, by definition, we have $C^1 = C = C^{\wedge_T^1}$.

Let $n \geq 2$ and assume that $C^{\wedge_T^{n-1}} = C^{n-1}$. By Proposition 2.17 and Lemma 2.16, we have

$$C^{\wedge_T^n} = C^{\wedge_T^{n-1}} \wedge_T C^{\wedge_T^1} = C^{n-1} \wedge_T C^1 = \text{Ker}(\tau_{n-1}) \wedge_T \text{Ker}(\tau_1) = \text{Ker}[(\tau_{n-1} \otimes \tau_1) \Delta_T]$$

so that

$$0 \rightarrow C^{\wedge_T^n} \hookrightarrow T^{(\tau_{n-1} \otimes \tau_1) \Delta_T} \left(\bigoplus_{a \geq n-1} M^{\square a} \right) \otimes \left(\bigoplus_{a \geq 1} M^{\square a} \right)$$

is an exact sequence. In order to conclude, it is enough to check that the following sequence

$$0 \rightarrow C^n \xrightarrow{\sigma_n} T^{(\tau_{n-1} \otimes \tau_1) \Delta_T} \left(\bigoplus_{a \geq n-1} M^{\square a} \right) \otimes \left(\bigoplus_{a \geq 1} M^{\square a} \right)$$

is exact. For every $0 \leq a \leq n-1$, we have

$$(\tau_{n-1} \otimes \tau_1) \Delta_T \sigma_n i_a^n = (\tau_{n-1} \otimes \tau_1) (\sigma_n \otimes \sigma_n) \Delta(n) i_a^n$$

By Proposition 2.8 we can write

$$\Delta(n) i_a^n = \sum_{r=0}^a (i_r^n \otimes i_{a-r}^n) f_{r,a-r}$$

where $f_{ij} : M^{\square a} \rightarrow M^{\square i} \otimes M^{\square j}$ are suitable morphisms. Thus we get

$$\begin{aligned} (\tau_{n-1} \otimes \tau_1) \Delta_T \sigma_n i_a^n &= \sum_{r=0}^a (\tau_{n-1} \otimes \tau_1) (\sigma_n \otimes \sigma_n) (i_r^n \otimes i_{a-r}^n) f_{r,a-r} \\ &= \sum_{r=0}^a (\tau_{n-1} i_r \otimes \tau_1 i_{a-r}) f_{r,a-r} = \sum_{r=0}^a (\lambda_r^{n-1} \otimes \lambda_{a-r}^1) f_{r,a-r} = 0 \end{aligned}$$

Therefore $(\tau_{n-1} \otimes \tau_1) \Delta_T \sigma_n = 0$, for every $n \in \mathbb{N}$.

Let $g : Y \rightarrow T$ be a morphism such that

$$(\tau_{n-1} \otimes \tau_1) \Delta_T g = 0.$$

Now, for every $c \in \mathbb{N}$ and for every $a \geq b$, we have

$$p_a \nu_b \tau_b i_c = p_a \nu_b \lambda_c^b = \begin{cases} p_a i_c & \text{for every } c \geq b \\ 0 = p_a i_c & \text{otherwise.} \end{cases}$$

so that

$$p_a \nu_b \tau_b = p_a, \text{ for every } a \geq b.$$

Thus, for every $a \geq n-1$ and $b \geq 1$, by Lemma 2.10, we have

$$0 = (p_a \nu_{n-1} \tau_{n-1} \otimes p_b \nu_1 \tau_1) \Delta_T g = (p_a \otimes p_b) \Delta_T g = \left(M^{\square a-1} \square_{\chi_M} M^{\square b} \right) p_{a+b} g.$$

By left exactness of the tensor functors, $M^{\square a-1} \square_{\chi_M} M^{\square b}$ is a monomorphism so that

$$p_{a+b} g = 0.$$

We conclude that

$$p_c g = 0, \text{ for every } c \geq n.$$

Set

$$\bar{g} = \pi_n g.$$

and let us prove that $g = \sigma_n \bar{g}$. By Lemma 2.14 this is the case if and only if

$$p_a g = p_a \sigma_n \bar{g}, \text{ for every } a \in \mathbb{N}.$$

We have

$$p_a \sigma_n \bar{g} = p_n^a \pi_n g = \begin{cases} p_a g & \text{for every } a < n \\ 0 = p_a g & \text{otherwise.} \end{cases}$$

□

3. TECHNICALITIES

The main aim of this section is to proof Theorem 3.11 that will be our main tool in the proof of Theorem 4.15.

LEMMA 3.1. *Let $\mathcal{M}$ be a monoidal category with left exact direct limits. Let $((A_i)_{i \in \mathbb{N}}, (\alpha_i^j)_{i,j \in \mathbb{N}})$ and $((B_i)_{i \in \mathbb{N}}, (\beta_i^j)_{i,j \in \mathbb{N}})$ be direct systems in $\mathcal{M}$, where, for $i \leq j$, $\alpha_i^j : A_i \rightarrow A_j$ and $\beta_i^j : B_i \rightarrow B_j$. Let $(\gamma_i : A_i \rightarrow B_i)_{i \in \mathbb{N}}$ be a direct system of monomorphisms. Let $(A, (\alpha_i)_{i \in \mathbb{N}}) = \varinjlim A_i$ and let $(\beta_i : B_i \rightarrow A)_{i \in \mathbb{N}}$ be a compatible family of monomorphisms such that $\beta_i \gamma_i = \alpha_i$ for any $i \in \mathbb{N}$. Then $(A, (\beta_i)_{i \in \mathbb{N}}) = \varinjlim B_i$.*

Proof. Since direct limits are left exact in $\mathcal{M}$, the canonical morphism $\varinjlim \beta_i : \varinjlim B_i \rightarrow A$ is a monomorphism. Moreover, since $\varinjlim \beta_i \circ \varinjlim \gamma_i = \varinjlim \alpha_i = \text{Id}_A$, we have that $\varinjlim \beta_i$ is also an epimorphism and hence an isomorphism. $\square$

3.2. Let $(\mathcal{M}, \otimes, \mathbf{1})$ be a cocomplete abelian monoidal category. Recall that a *graded* coalgebra in $\mathcal{M}$ is a coalgebra (B, Δ, ε) endowed with a family (B_i, β_i) of subobjects of B , such that

$$B = \oplus_{i \in \mathbb{N}} B_i$$

and there exists a family $(\Delta_i)_{i \in \mathbb{N}}$ of morphisms

$$\Delta_i : B_i \rightarrow (B \otimes B)_i = \oplus_{a+b=i} (B_a \otimes B_b),$$

such that

$$\Delta \beta_i = [\nabla_{a+b=i} (\beta_a \otimes \beta_b)] \Delta_i$$

and

$$\varepsilon \beta_i = 0, \text{ for every } i \geq 1.$$

Here $\nabla_{a+b=i} (\beta_a \otimes \beta_b)$ denotes the codiagonal morphism associated to the family $(\beta_a \otimes \beta_b)_{a+b=i}$. In particular it follows that ε restricts to a morphism

$$\varepsilon_0 : B_0 \rightarrow \mathbf{1}$$

such that

$$\varepsilon = \varepsilon_0 \beta_0$$

and $(B_0, \Delta_0, \varepsilon_0)$ is a coalgebra in $\mathcal{M}$. Moreover β_0 is a coalgebra homomorphism.

PROPOSITION 3.3. *Let $(\mathcal{M}, \otimes, \mathbf{1})$ be a cocomplete abelian monoidal category. Let $B = \oplus_{i \in \mathbb{N}} B_i$ be a graded coalgebra. Denote by (L, p) the cokernel of β_0 in $\mathcal{M}$. Then*

$$(23) \quad p^{\otimes n+1} \Delta_B^n \beta_b = 0, \text{ for every } 0 \leq b \leq n.$$

Moreover

$$B = \varinjlim (B_0^{\wedge^i_B})_{i \in \mathbb{N}}.$$

Proof. Denote by $\beta_i : B_i \rightarrow B$ the canonical inclusion and denote by $\tau_i : B \rightarrow B_i$ the canonical projection, for every $i \in \mathbb{N}$. Since β_0 is a coalgebra homomorphism and β_0 is a monomorphism, we can consider

$$(B_0^{\wedge^n_B}, \delta_n) := \text{Ker}(p^{\otimes n} \Delta_B^{n-1}).$$

Denote by $\xi_i^j : B_0^{\wedge^i_B} \rightarrow B_0^{\wedge^j_B}$ the canonical inclusion, for every $j \geq i$. Hence we have

$$\delta_j \xi_i^j = \delta_i.$$

In order to prove (23), we proceed by induction on $n \geq 0$. For $n = 0$, then $b = 0$ and we have

$$p^{\otimes n+1} \Delta_B^n \beta_b = p \Delta_B^0 \beta_0 = p \beta_0 = 0.$$

Let $n \geq 1$ and assume $p^{\otimes i+1} \Delta_B^i \beta_j = 0$, for every $0 \leq j \leq i \leq n-1$. For every $0 \leq c \leq n$, we have

$$\begin{aligned} p^{\otimes n+1} \Delta_B^n \beta_c &= (p^{\otimes n-1} \otimes p^{\otimes 2}) (\Delta_B^{n-2} \otimes \Delta_B) \Delta_B \beta_c \\ &= (p^{\otimes n-1} \Delta_B^{n-2} \otimes p^{\otimes 2} \Delta_B) [\nabla_{a+b=c} (\beta_a \otimes \beta_b)] \Delta_c \\ &= [\nabla_{a+b=c} (p^{\otimes n-1} \Delta_B^{n-2} \beta_a \otimes p^{\otimes 2} \Delta_B \beta_b)] \Delta_c = 0. \end{aligned}$$

By definition of $(B_0^{\wedge_B^n}, \delta_n)$, there exists a unique morphism

$$\gamma_n : \oplus_{i=0}^n B_i \rightarrow B_0^{\wedge_B^{n+1}}$$

such that

$$\delta_{n+1}\gamma_n = \nabla_{i=0}^n \beta_i.$$

Since each β_i cosplits, then $\nabla_{i=0}^n \beta_i$ is a monomorphism. Thus also γ_n is a monomorphism. Denote by $\beta_a^b : \oplus_{i=0}^a B_i \rightarrow \oplus_{i=0}^b B_i$ the canonical injection when $a \leq b$. Then we have

$$\delta_{n+2}\gamma_{n+1}\beta_n^{n+1} = (\nabla_{i=0}^{n+1} \beta_i) \beta_n^{n+1} = \nabla_{i=0}^n \beta_i = \delta_{n+1}\gamma_n = \delta_{n+2}\xi_{n+1}^{n+2}\gamma_n$$

Since δ_{n+2} is a monomorphism, we get that

$$\gamma_{n+1}\beta_n^{n+1} = \xi_{n+1}^{n+2}\gamma_n$$

for every $n \in \mathbb{N}$. Thus $(\gamma_n : \oplus_{i=0}^n B_i \rightarrow B_0^{\wedge_B^{n+1}})_{n \in \mathbb{N}}$ defines a direct system of monomorphisms in $\mathcal{M}$. Now, as, by Proposition 2.5, $(B, (\nabla_{i=0}^n \beta_i)_{n \in \mathbb{N}}) = \varinjlim (\oplus_{i=0}^n B_i)$, by Lemma 3.1 we have that

$$(B, (\delta_n)_{n \in \mathbb{N}}) = \varinjlim (B_0^{\wedge_B^i})_{i \in \mathbb{N}}. \quad \square$$

PROPOSITION 3.4. *Let $i \in \{1, 2\}$. Let $f_i : X_i \rightarrow Y_i$ be morphisms in an abelian monoidal category $\mathcal{M}$. Let $\sigma_i : Y_i \rightarrow X_i$ such that $f_i \sigma_i = \text{Id}_{Y_i}$. Then*

$$\text{Ker}(f_1 \otimes f_2) = [\text{Ker}(f_1) \otimes X_2] + [X_1 \otimes \text{Ker}(f_2)].$$

Proof. Let $(K_i, k_i) = \text{Ker}(f_i)$ for $i = 1, 2$. Let $\nu_1 : K_1 \otimes X_2 \rightarrow (K_1 \otimes X_2) \oplus (X_1 \otimes K_2)$ and $\nu_2 : X_1 \otimes K_2 \rightarrow (K_1 \otimes X_2) \oplus (X_1 \otimes K_2)$ be the canonical inclusions. Then, by the universal property of coproducts, there is a unique morphism $\tau : (K_1 \otimes X_2) \oplus (X_1 \otimes K_2) \rightarrow X_1 \otimes X_2$ such that

$$(24) \quad \tau \nu_1 = k_1 \otimes X_2 \quad \text{and} \quad \tau \nu_2 = X_1 \otimes k_2.$$

By definition, one has $(K_1 \otimes X_2) + (X_1 \otimes K_2) = \text{Im}(\tau) = \text{Ker}(\pi)$, where $(C, \pi) = \text{coker}(\tau)$.

Thus, in order to prove our statement, we will show that $(C, \pi) = (Y_1 \otimes Y_2, f_1 \otimes f_2)$. By (24), we have

$$\begin{aligned} (f_1 \otimes f_2)\tau \nu_1 &= (f_1 \otimes f_2)(k_1 \otimes X_2) = 0, \\ (f_1 \otimes f_2)\tau \nu_2 &= (f_1 \otimes f_2)(X_1 \otimes k_2) = 0, \end{aligned}$$

so that $(f_1 \otimes f_2)\tau = 0$. By the universal property of cokernels, we obtain a unique morphism $\alpha : C \rightarrow Y_1 \otimes Y_2$ such that $\alpha\pi = f_1 \otimes f_2$.

Define $\beta : Y_1 \otimes Y_2 \rightarrow C$ by $\beta := \pi(\sigma_1 \otimes \sigma_2)$. Let us prove that β is a two-sided inverse of α . Clearly one has

$$\alpha\beta = \alpha\pi(\sigma_1 \otimes \sigma_2) = (f_1 \otimes f_2)(\sigma_1 \otimes \sigma_2) = \text{Id}_{X_1 \otimes X_2}.$$

Now, since $f_i \sigma_i = \text{Id}_{Y_i}$, there is a unique morphism $\rho_i : X_i \rightarrow K_i$ such that $\rho_i k_i = \text{Id}_{K_i}$ and

$$(25) \quad k_i \rho_i + \sigma_i f_i = \text{Id}_{X_i}, \text{ for any } i \in \{1, 2\}.$$

Then we have:

$$\begin{aligned} \beta\alpha\pi &= \beta(f_1 \otimes f_2) \\ &= \pi(\sigma_1 f_1 \otimes \sigma_2 f_2) \\ (25) \quad &\stackrel{=}{=} \pi[\sigma_1 f_1 \otimes (\text{Id}_{X_2} - k_2 \rho_2)] \\ &= \pi(\sigma_1 f_1 \otimes \text{Id}_{X_2}) - \pi(\sigma_1 f_1 \otimes k_2 \rho_2) \\ &= \pi[(\text{Id}_{X_1} - k_1 \rho_1) \otimes \text{Id}_{X_2}] - \pi(X_1 \otimes k_2)(\sigma_1 f_1 \otimes \rho_2) \\ (24) \quad &\stackrel{=}{=} \pi - \pi(k_1 \rho_1 \otimes \text{Id}_{X_2}) - \pi\tau \nu_2(\sigma_1 f_1 \otimes \rho_2) \\ &= \pi - \pi(k_1 \otimes \text{Id}_{X_2})(\rho_1 \otimes \text{Id}_{X_2}) \\ (24) \quad &\stackrel{=}{=} \pi - \pi\tau \nu_1(\rho_1 \otimes \text{Id}_{X_2}) \\ &= \pi \end{aligned}$$

Since π is an epimorphism we conclude that $\beta\alpha = \text{Id}_C$ and hence that α is an isomorphism. Thus $(C, \pi) = (Y_1 \otimes Y_2, f_1 \otimes f_2)$. $\square$

PROPOSITION 3.5. *Let $\delta : D \rightarrow C$ be a morphism that cosplits in $\mathcal{M}$. If δ is a coalgebra homomorphism, then we have*

$$(26) \quad D^{\wedge_C^2} = D \wedge_C D = \Delta_C^{-1}(D \otimes C + C \otimes D).$$

Proof. Set $(L, p) = \text{coker}(\sigma)$. Let

$$(D^{\wedge_C^n}, \delta_n) := \text{Ker}(p^{\otimes n} \Delta_C^{n-1}).$$

We have

$$\begin{aligned} D^{\wedge_C^2} &= \text{Ker}[(p \otimes p) \Delta_C] \\ &\stackrel{(*)}{=} \Delta_C^{-1}[\text{Ker}(p \otimes p)] \\ &\stackrel{(**)}{=} \Delta_C^{-1}\{[\text{Ker}(p) \otimes C] + [C \otimes \text{Ker}(p)]\} = \Delta_C^{-1}[(D \otimes C) + (C \otimes D)]. \end{aligned}$$

where in $(*)$ we have applied ([St, Proposition 5.1, page 90]) and in $(**)$ Proposition 3.4. $\square$

LEMMA 3.6. *Let $\sigma : A \rightarrow A'$ be a coalgebra homomorphism in an abelian monoidal category $\mathcal{M}$. Assume that σ is a monomorphism. Let $i \in \{1, 2\}$ and let $\alpha_i : X_i \rightarrow A$, $\alpha'_i : X'_i \rightarrow A$, $\sigma_i : X_i \rightarrow X'_i$ be morphism such that $\sigma\alpha_i = \alpha'_i\sigma_i$. Let $(L_i, p_i) = \text{coker}(\alpha_i)$ and $(L'_i, p'_i) = \text{coker}(\alpha'_i)$ and let*

$$(X_1 \wedge_A X_2, \lambda_{1,2}) = \text{Ker}[(p_1 \otimes p_2) \Delta_A] \quad \text{and} \quad (X'_1 \wedge_{A'} X'_2, \lambda'_{1,2}) = \text{Ker}[(p'_1 \otimes p'_2) \Delta_{A'}].$$

Then there is a (unique) morphism

$$\lambda = \lambda(A, X_1, X_2; A', X'_1, X'_2) : X_1 \wedge_A X_2 \rightarrow X'_1 \wedge_{A'} X'_2$$

such that $\lambda'_{1,2}\lambda = \sigma\lambda_{1,2}$. Moreover λ is a monomorphism.

Proof. Since

$$p'_i\sigma\alpha_i = p'_i\alpha'_i\sigma_i = 0,$$

by the universal property of cokernels, there is a unique morphism $\tau_i : L_i \rightarrow L'_i$ such that $\tau_i p_i = p'_i \sigma$. Then we have:

$$(p'_1 \otimes p'_2) \Delta_{A'} \sigma \lambda_{1,2} = (p'_1 \otimes p'_2) (\sigma \otimes \sigma) \Delta_A \lambda_{1,2} = (\tau_1 \otimes \tau_2) (p_1 \otimes p_2) \Delta_A \lambda_{1,2} = 0.$$

By the universal property of kernels, there is a unique morphism $\lambda : X_1 \wedge_A X_2 \rightarrow X'_1 \wedge_{A'} X'_2$ such that $\lambda'_{1,2}\lambda = \sigma\lambda_{1,2}$. Clearly, as σ and $\lambda_{1,2}$ are monomorphisms, λ is a monomorphism too. $\square$

LEMMA 3.7. *Let $((X_i)_{i \in \mathbb{N}}, (\xi_i^j)_{i,j \in \mathbb{N}})$ and let $((Y_i)_{i \in \mathbb{N}}, (\zeta_i^j)_{i,j \in \mathbb{N}})$ be direct systems in a monoidal category $\mathcal{M}$, where, for $i \leq j$, $\xi_i^j : X_i \rightarrow X_j$ and $\zeta_i^j : Y_i \rightarrow Y_j$. Let $\sigma : A \rightarrow B$ be a coalgebra homomorphism and let $(\alpha_i : X_i \rightarrow A)_{i \in \mathbb{N}}$ and $(\beta_i : Y_i \rightarrow B)_{i \in \mathbb{N}}$ be compatible families of morphisms in $\mathcal{M}$. Let $\lambda_i : X_i \rightarrow Y_i$ be a morphism such that $\beta_i \lambda_i = \sigma \alpha_i$, for any $i \in \mathbb{N}$. If β_i is a monomorphism, for any $i \in \mathbb{N}$, we have that $(\lambda_i : X_i \rightarrow Y_i)_{i \in \mathbb{N}}$ is a direct system of morphisms in $\mathcal{M}$.*

Proof. For any $i \leq j$, we have that:

$$\beta_j \lambda_j \xi_i^j = \sigma \alpha_j \xi_i^j = \sigma \alpha_i = \beta_i \lambda_i = \beta_j \zeta_i^j \lambda_i.$$

Since β_j is a monomorphism for any $j \in \mathbb{N}$, we conclude that $\lambda_j \xi_i^j = \zeta_i^j \lambda_i$ i.e. that $(\lambda_i : X_i \rightarrow Y_i)_{i \in \mathbb{N}}$ is a direct system of morphisms in $\mathcal{M}$. $\square$

LEMMA 3.8. *Let $\mathcal{M}$ be a cocomplete monoidal category with left exact direct limits. Let $((X_i)_{i \in \mathbb{N}}, (\xi_i^j)_{i,j \in \mathbb{N}})$ be a direct system in $\mathcal{M}$, where, for $i \leq j$, $\xi_i^j : X_i \rightarrow X_j$.*

Let $\gamma : \mathbb{N} \rightarrow \mathbb{N}$ be an injection. Then $((X_{\gamma(i)})_{i \in \mathbb{N}}, (\xi_{\gamma(i)}^{\gamma(j)})_{i,j \in \mathbb{N}})$ is a direct system in $\mathcal{M}$. Let $(X, (\lambda_i)_{i \in \mathbb{N}}) = \varinjlim X_{\gamma(i)}$, where $\lambda_i : X_{\gamma(i)} \rightarrow X$ for any $i \in \mathbb{N}$.

Then $(X, (\xi_i)_{i \in \mathbb{N}}) = \varinjlim X_i$, where $\xi_i : X_i \rightarrow X$ is defined by $\xi_i := \lambda_j \xi_i^{\gamma(j)} : X_i \rightarrow X$, where $j \in \mathbb{N}$ is such that $\gamma(j) \geq i$.

Proof. Clearly $((X_{\gamma(i)})_{i \in \mathbb{N}}, (\xi_{\gamma(i)}^{\gamma(j)})_{i, j \in \mathbb{N}})$ is a direct system. Let us prove the last assertion. Let $j, j' \in \mathbb{N}$ such that $\gamma(j') \geq \gamma(j) \geq i$. Then

$$\lambda_{j'} \xi_i^{\gamma(j')} = \lambda_{j'} \xi_{\gamma(j)}^{\gamma(j')} \xi_i^{\gamma(j)} = \lambda_j \xi_i^{\gamma(j)},$$

so that ξ_i is well defined. Note that

$$(27) \quad \xi_{\gamma(j)} = \lambda_j \xi_{\gamma(j)}^{\gamma(j)} = \lambda_j.$$

Moreover, for any $i \leq j$, and for any $t \in \mathbb{N}$ such that $\gamma(t) \geq j$ we have:

$$\xi_j \xi_i^j = \lambda_t \xi_j^{\gamma(t)} \xi_i^j = \lambda_t \xi_i^{\gamma(t)} = \xi_i,$$

so that $(\xi_i : X_i \rightarrow X)_{i \in \mathbb{N}}$ is a direct system of morphisms. Let now $(f_i : X_i \rightarrow Y)_{i \in \mathbb{N}}$ be a compatible family of morphisms in $\mathcal{M}$. Then $(f_{\gamma(i)} : X_{\gamma(i)} \rightarrow Y)_{i \in \mathbb{N}}$ is a compatible family of morphisms in $\mathcal{M}$ so that there exists a unique morphism $f : X \rightarrow Y$ such that $f \lambda_i = f_{\gamma(i)}$ for any $i \in \mathbb{N}$. For any $i \in \mathbb{N}$ and for any $j \in \mathbb{N}$ is such that $\gamma(j) \geq i$, we obtain

$$f \xi_i = f \lambda_j \xi_i^{\gamma(j)} = f_{\gamma(i)} \xi_i^{\gamma(j)} = f_i.$$

Let $g : X \rightarrow Y$ be another morphism such that $g \xi_i = f_i$. Then we have

$$g \lambda_i \stackrel{(27)}{=} g \xi_{\gamma(i)} = f_{\gamma(i)}.$$

By uniqueness of f we get $g = f$, so that $(X, (\xi_i)_{i \in \mathbb{N}}) = \varinjlim X_i$. $\square$

LEMMA 3.9. *Let $(\mathcal{M}, \otimes, \mathbf{1})$ be a cocomplete abelian monoidal category satisfying AB5, with left exact direct limits and left and right exact tensor functors. Let $(\mathbb{I}, \leq)$ be a directed partially ordered set. Let $((X_i)_{i \in \mathbb{I}}, (\xi_i^j)_{i, j \in \mathbb{I}})$ be a direct system in $\mathcal{M}$, where, for $i \leq j$, $\xi_i^j : X_i \rightarrow X_j$. Let $(w_i : X_i \rightarrow W)_{i \in \mathbb{I}}$ be a compatible family of monomorphisms in $\mathcal{M}$. Let $(X, (\xi_i)_{i \in \mathbb{I}}) = \varinjlim X_i$. Let $w : X \rightarrow W$ be the unique morphism such that $w \xi_i = w_i$ for every i . Let $\xi : \oplus X_i \rightarrow X$ be the unique morphism such that $\xi \varepsilon_i = \xi_i$ for any $i \in \mathbb{I}$ and let $\omega : \oplus X_i \rightarrow W$ be the unique morphism such that $\omega \varepsilon_i = w_i$ for any $i \in \mathbb{I}$, where $\varepsilon_i : X_i \rightarrow \oplus X_i$ is the canonical inclusion. Then:*

$$w \xi = \omega.$$

Moreover w is a monomorphism and ξ is an epimorphism.

Proof. Since $w \xi_i = w_i$, the ξ_i 's are monomorphisms. Clearly we have

$$w \xi \varepsilon_i = w \xi_i = w_i = \omega \varepsilon_i \text{ for any } i \in \mathbb{I}$$

and hence

$$w \xi = \omega$$

Moreover, regarding $(w_i : X_i \rightarrow W)_{i \in \mathbb{I}}$ as a direct system of monomorphism, in view of AB5, we have that w is a monomorphism and ξ is an epimorphism. $\square$

LEMMA 3.10. *Let $(\mathcal{M}, \otimes, \mathbf{1})$ be a cocomplete abelian monoidal category satisfying AB5 and with left and right exact tensor functors. Let $(\mathbb{I}, \leq)$ be a directed partially ordered set. Let $((X_i)_{i \in \mathbb{I}}, (\xi_i^j)_{i, j \in \mathbb{I}})$ be a direct system in $\mathcal{M}$, where, for $i \leq j$, $\xi_i^j : X_i \rightarrow X_j$. If $\oplus X_i$ commutes with $\otimes$, then $\varinjlim X_i$ does.*

Proof. Let $((X_i)_{i \in \mathbb{I}}, (\xi_i^j)_{i, j \in \mathbb{I}})$ be a direct system in $\mathcal{M}$, where, for $i \leq j$, $\xi_i^j : X_i \rightarrow X_j$. Let $(X, (\xi_i)_{i \in \mathbb{I}}) = \varinjlim X_i$ and let Y be an object in $\mathcal{M}$. By [St, Lemma 1.2, page 115], the ξ_i 's are monomorphisms. Also, by the universal property of the coproduct, there is a unique morphism $\xi : \oplus X_i \rightarrow X$ such that

$$(28) \quad \xi \varepsilon_i = \xi_i \text{ for any } i \in \mathbb{I}$$

where $\varepsilon_i : X_i \rightarrow \oplus X_i$ is the canonical inclusion. Moreover, ξ is an epimorphism.

Assume that

$$(29) \quad ((\oplus X_i) \otimes Y, \varepsilon_i \otimes Y) = \oplus (X_i \otimes Y),$$

Let $\gamma_i : X_i \otimes Y \rightarrow \varinjlim (X_i \otimes Y)$ be the canonical morphism. By the universal property of coproduct and by (29), there is a unique morphism $\gamma : (\oplus X_i) \otimes Y \rightarrow \varinjlim (X_i \otimes Y)$ such that

$$(30) \quad \gamma(\varepsilon_i \otimes Y) = \gamma_i \text{ for any } i \in \mathbb{I}.$$

In an analogous way, by the universal property of direct limits, there is a unique morphism $\Lambda : \varinjlim (X_i \otimes Y) \rightarrow X \otimes Y$ such that

$$(31) \quad \Lambda \gamma_i = \xi_i \otimes Y \text{ for any } i \in \mathbb{I}.$$

It is easy to see that we can apply Lemma 3.9 to the present situation and get:

$$\Lambda \gamma = \xi \otimes Y.$$

where Λ is a monomorphism and γ is an epimorphism. Moreover, since the tensor functor is left exact and ξ is an epimorphism, we get that $\xi \otimes Y$ is an epimorphism. Hence Λ is an epimorphism too. $\square$

THEOREM 3.11. *Let (C, Δ, ε) be a coalgebra in a cocomplete abelian monoidal category $\mathcal{M}$ satisfying AB5, with left and right exact tensor functors. Assume that denumerable coproducts commute with $\otimes$. Let $\alpha : C \rightarrow A$ and $\sigma : A \rightarrow B$ be monomorphisms which are coalgebra homomorphisms and let $\beta = \sigma\alpha$. Assume that σ cosplits in $\mathcal{M}$. Let $p_\alpha = \text{coker}(\alpha)$ in $\mathcal{M}$, let $(C^{\wedge_A^n}, \alpha_n) := \text{Ker}(p_\alpha^{\otimes n} \Delta_A^{n-1})$ and assume that α_n cosplits in $\mathcal{M}$, for every $n \in \mathbb{N}$. If $\tilde{C}_A = A$ and $B = A^{\wedge_B^2}$, then $\tilde{C}_B = B$.*

Proof. For any morphism η we set $(L_\eta, p_\eta) = \text{coker}(\eta)$ in $\mathcal{M}$. By Proposition 3.5, we get

$$B = A^{\wedge_B^2} \stackrel{(26)}{=} \Delta_B^{-1}(A \otimes B + B \otimes A).$$

Let

$$(C^{\wedge_A^n}, \alpha_n) := \text{Ker}(p_\alpha^{\otimes n} \Delta_A^{n-1}) \quad \text{and} \quad (C^{\wedge_B^n}, \beta_n) := \text{Ker}(p_\beta^{\otimes n} \Delta_B^{n-1}).$$

By assumption $(A, (\alpha_n)_{n \in \mathbb{N}}) = \tilde{C}_A = \varinjlim C^{\wedge_A^n}$. Then, by Lemma 3.10, we obtain:

$$A \otimes B = (\varinjlim C^{\wedge_A^n}) \otimes B = \varinjlim (C^{\wedge_A^n} \otimes B).$$

We have:

$$\begin{aligned} B &= \Delta_B^{-1}[\varinjlim (C^{\wedge_A^n} \otimes B) + \varinjlim (B \otimes C^{\wedge_A^n})] \\ &= \Delta_B^{-1} \varinjlim [(C^{\wedge_A^n} \otimes B) + (B \otimes C^{\wedge_A^n})] \\ &= \varinjlim \Delta_B^{-1} [(C^{\wedge_A^n} \otimes B) + (B \otimes C^{\wedge_A^n})] \\ &= \varinjlim (C^{\wedge_A^n} \wedge_B C^{\wedge_A^n}) \end{aligned}$$

where in the second equality we have used that in an AB5-category direct limits of direct systems of subobjects are just sums of their respective families; in the third we have used a well known property of AB5-categories (see [St, Proposition 1.1, page 114]); in the last equality we have used Proposition 3.5 in the case $\delta = \sigma\alpha_n : C^{\wedge_A^n} \rightarrow B$. Note that, by applying inductively Lemma 3.6, we obtain, for any $n \in \mathbb{N}$, a monomorphism

$$\lambda_{A,B}^n : C^{\wedge_A^n} \rightarrow C^{\wedge_B^n}$$

such that

$$(32) \quad \beta_n \lambda_{A,B}^n = \sigma \alpha_n.$$

By Lemma 3.7, $(\lambda_{A,B}^n : C^{\wedge_A^n} \rightarrow C^{\wedge_B^n})_{i \in \mathbb{N}}$ is a direct system of monomorphisms in $\mathcal{M}$. Let $m \leq n$. Note that, if we denote by $\xi_{A,m}^n : C^{\wedge_A^m} \rightarrow C^{\wedge_A^n}$ and by $\xi_{B,m}^n : C^{\wedge_B^m} \rightarrow C^{\wedge_B^n}$ the canonical morphisms, this means that $\lambda_{A,B}^n \xi_{A,m}^n = \xi_{B,m}^n \lambda_{A,B}^m$.

Let $(L_n, p_n) = \text{coker}(\sigma_n \alpha_n)$ and $(L'_n, p'_n) = \text{coker}(\beta_n)$ and let

$$(C^{\wedge_A^n} \wedge_B C^{\wedge_A^n}, \lambda_n) = \text{Ker}[(p_n \otimes p_n) \Delta_B] \quad \text{and} \quad (C^{\wedge_B^n} \wedge_B C^{\wedge_B^n}, \lambda'_n) = \text{Ker}[(p'_n \otimes p'_n) \Delta_B].$$

In view of (32), we can apply Lemma 3.6 to obtain a monomorphism

$$\lambda(n) = \lambda(B, C^{\wedge_A^n}, C^{\wedge_A^n}; B, C^{\wedge_B^n}, C^{\wedge_B^n}) : C^{\wedge_A^n} \wedge_B C^{\wedge_A^n} \rightarrow C^{\wedge_B^n} \wedge_B C^{\wedge_B^n}$$

such that

$$\lambda'_n \lambda(n) = \lambda_n.$$

By Lemma 3.7, $(\lambda(n) : C^{\wedge_A^n} \wedge_B C^{\wedge_A^n} \rightarrow C^{\wedge_B^n} \wedge_B C^{\wedge_B^n})_{n \in \mathbb{N}}$ is a direct system of monomorphisms in $\mathcal{M}$. Since, by the foregoing, $B = \varinjlim (C^{\wedge_A^n} \wedge_B C^{\wedge_A^n})$, by Lemma 3.1, applied in the case $\gamma_i = \lambda(i)$ for any $i \in \mathbb{N}$, we obtain that

$$(B, (\lambda'_n)_{n \in \mathbb{N}}) = \varinjlim (C^{\wedge_B^n} \wedge_B C^{\wedge_B^n}).$$

As

$$(C^{\wedge_B^n} \wedge_B C^{\wedge_B^n}, \lambda'_n) \stackrel{(22)}{=} (C^{\wedge_B^{2n}}, \beta_{2n})$$

we obtain

$$(B, (\beta_{2n})_{n \in \mathbb{N}}) = \varinjlim C^{\wedge_B^{2n}}.$$

Now, apply Lemma 3.8 in the case when $\gamma : \mathbb{N} \rightarrow \mathbb{N}$ is defined by setting $\gamma(n) = 2n$ for every $n \in \mathbb{N}$. Then we get

$$(B, (\beta_n)_{n \in \mathbb{N}}) = \varinjlim C^{\wedge_B^{2n}} = \tilde{C}_B.$$

□

4. FORMAL SMOOTHNESS OF THE COTENSOR COALGEBRA

The main aim of this section is to prove Theorem 4.15 which asserts that the cotensor coalgebra $T_C^c(M)$ is formally smooth whenever C is a formally smooth coalgebra in a cocomplete and complete abelian monoidal category $\mathcal{M}$ satisfying $AB5$, and M is an $\mathcal{I}$ -injective C -bicomodule in $\mathcal{M}$.

DEFINITION 4.1. Let (C, Δ, ε) be a coalgebra in $(\mathcal{M}, \otimes, \mathbf{1})$ and let (L, ρ_L^l, ρ_L^r) be a C -bicomodule. Let us consider *the standard complex*:

$$0 \longrightarrow \mathcal{M}(L, \mathbf{1}) \xrightarrow{b^0} \mathcal{M}(L, C) \xrightarrow{b^1} \mathcal{M}(L, C \otimes C) \xrightarrow{b^2} \mathcal{M}(L, C \otimes C \otimes C) \xrightarrow{b^3} \dots$$

Let $n \in \mathbb{N}$. For every $0 \leq i \leq n+1$ and for every $f \in \mathcal{M}(L, C^{\otimes n})$, we define, for $n = 0$:

$$b_0^0 = l_C \circ (f \otimes C) \circ \rho_L^r, \quad b_1^0 = r_C \circ (C \otimes f) \circ \rho_L^l$$

and, for $n > 0$:

$$b_i^n(f) = \begin{cases} (f \otimes C) \circ \rho_L^r & i = 0; \\ (C^{\otimes n-i} \otimes \Delta \otimes C^{\otimes i-1}) \circ f, & i = 1, \dots, n; \\ (C \otimes f) \circ \rho_L^l & i = n+1. \end{cases}$$

One has that

$$b^n(f) = \sum_{i=0}^{n+1} (-1)^i b_i^n(f), \text{ for every } n \geq 0.$$

In particular, for $n \in \{0, 1, 2\}$ the differentials b^n are given by:

$$\begin{aligned} b^0(f) &= l_C \circ (f \otimes C) \circ \rho_L^r - r_C \circ (C \otimes f) \circ \rho_L^l; \\ b^1(f) &= (f \otimes C) \circ \rho_L^r - \Delta \circ f + (C \otimes f) \circ \rho_L^l; \\ b^2(f) &= (f \otimes C) \circ \rho_L^r - (C \otimes \Delta) \circ f + (\Delta \otimes C) \circ f - (C \otimes f) \circ \rho_L^l. \end{aligned}$$

Further details can be found in [AMS].

4.2. Let $\mathcal{M}$ be an abelian category and let $\mathcal{H}$ be a class of monomorphisms in $\mathcal{M}$. We recall that an object I in $\mathcal{M}$ is called *injective rel* λ , where $\lambda : X \rightarrow Y$ is a monomorphism in $\mathcal{H}$, if $\mathcal{M}(\lambda, I) : \mathcal{M}(Y, I) \rightarrow \mathcal{M}(X, I)$ is surjective. I is called $\mathcal{H}$ -injective if it is injective rel λ for every λ in $\mathcal{H}$. The *closure* of $\mathcal{H}$ is the class $\mathcal{C}(\mathcal{H})$ containing all monomorphisms λ in $\mathcal{M}$ such that every $\mathcal{H}$ -injective object is also injective rel λ . The class $\mathcal{H}$ is called *closed* if $\mathcal{H} = \mathcal{C}(\mathcal{H})$. A closed class $\mathcal{H}$ is called *injective* if for any object M in $\mathcal{M}$ there is a monomorphism $\lambda : M \rightarrow I$ in $\mathcal{H}$ such that I is $\mathcal{H}$ -injective.

4.3. We fix a coalgebra C in an abelian monoidal category $\mathcal{M}$. Let $\mathbb{U} : {}^C\mathcal{M}^C \rightarrow \mathcal{M}$ be the forgetful functor. Then

$$(33) \quad \mathcal{I} := \{f \in {}^C\mathcal{M}^C \mid \mathbb{U}(f) \text{ cosplits in } \mathcal{M}\}.$$

is an injective class of monomorphisms.

Now, for any C -bicomodule $M \in {}^C\mathcal{M}^C$, we define the Hochschild cohomology of C with coefficients in M by:

$$\mathbf{H}^\bullet(M, C) = \mathbf{Ext}_{\mathcal{I}}^\bullet(M, C),$$

where $\mathbf{Ext}_{\mathcal{I}}^\bullet(M, -)$ are the relative left derived functors of ${}^C\mathcal{M}^C(M, -)$. The notion of Hochschild cohomology for algebras and coalgebras in monoidal categories has been deeply investigated in [AMS]. Here we quote some results that will be needed afterwards.

DEFINITION 4.4. A coalgebra in $\mathcal{M}$ is called *coseparable* if the comultiplication $\Delta : C \rightarrow C \otimes C$ has a retraction in ${}^C\mathcal{M}^C$.

THEOREM 4.5. [AMS, Theorem 4.4] *Let C be a coalgebra in an abelian monoidal category $\mathcal{M}$. Then the following assertions are equivalent:*

- (a) C is coseparable.
- (b) C is $\mathcal{I}$ -injective.
- (c) $\mathbf{H}^1(M, C) = 0$, for all $M \in {}^C\mathcal{M}^C$.
- (d) $\mathbf{H}^n(M, C) = 0$, for all $M \in {}^C\mathcal{M}^C$ and $n > 0$.
- (e) Any morphism in ${}^C\mathcal{M}^C$ cosplits in ${}^C\mathcal{M}^C$ whenever it cosplits in $\mathcal{M}$.
- (f) The category ${}^C\mathcal{M}^C$ is $\mathcal{I}$ -cosemisimple (i.e. every object in ${}^C\mathcal{M}^C$ is $\mathcal{I}$ -injective).

DEFINITION 4.6. An extension $\sigma : C \rightarrow E$ is a *trivial extension* whenever it admits a retraction that is a coalgebra homomorphism.

DEFINITION 4.7. We say that the sequence $\left((E_i)_{i \in \mathbb{N}}, (\eta_i^j)_{i, j \in \mathbb{N}}\right)$ of morphisms in an abelian monoidal category $\mathcal{M}$

$$(34) \quad E_1 \xrightarrow{\eta_1^2} E_2 \xrightarrow{\eta_2^3} \dots \xrightarrow{\eta_{n-1}^n} E_n \xrightarrow{\eta_n^{n+1}} E_{n+1} \xrightarrow{\eta_{n+1}^{n+2}} \dots$$

is a *direct system of extensions* if η_n^{n+1} is a coalgebra homomorphism and $E_n \wedge_{E_{n+1}} E_n = E_{n+1}$, for any $n \geq 1$. We say that a direct system of extensions $\left((E_i)_{i \in \mathbb{N}}, (\eta_i^j)_{i, j \in \mathbb{N}}\right)$ is a *direct system of Hochschild extensions* if each η_n^{n+1} has a retraction in $\mathcal{M}$.

EXAMPLE 4.8. Let C and E be coalgebras in an abelian monoidal category $\mathcal{M}$. Let $\delta : C \rightarrow E$ be a monomorphism which is a homomorphism of coalgebras in $\mathcal{M}$. The sequence $\left((C^{\wedge_E^i})_{i \in \mathbb{N}}, (\xi_i^j)_{i, j \in \mathbb{N}}\right)$ is a direct system of coalgebras extensions that will be called the *C -adic direct system in E* .

DEFINITION 4.9. Let $\mathcal{M}$ be an abelian monoidal category. We say that the direct system of extensions $\left((E_i)_{i \in \mathbb{N}}, (\eta_i^j)_{i, j \in \mathbb{N}}\right)$ has a *direct limit* if $\varinjlim E_i$ exists in the category $\mathbf{Coalg}(\mathcal{M})$ of coalgebras in $\mathcal{M}$.

REMARK 4.10. If $\left((E_i)_{i \in \mathbb{N}}, (\eta_i^j)_{i, j \in \mathbb{N}}\right)$ is a direct system of Hochschild extensions, then, for any $n \geq 1$,

$$0 \longrightarrow E_n \xrightarrow{\eta_n^{n+1}} E_{n+1} \longrightarrow \mathrm{Coker}(\eta_n^{n+1}) \longrightarrow 0$$

is a Hochschild extension of E_n with cokernel $\mathrm{Coker}(\eta_n^{n+1})$.

THEOREM 4.11. [AMS, Theorem 4.16] *Let (C, Δ, ε) be a coalgebra in $\mathcal{M}$. Then the following conditions are equivalent:*

- (a) *The canonical map $\mathrm{Hom}_{\mathbf{Coalg}}(\delta, C) : \mathrm{Hom}_{\mathbf{Coalg}}(E, C) \rightarrow \mathrm{Hom}_{\mathbf{Coalg}}(D, C)$ is surjective for every coalgebra homomorphism $\delta : D \rightarrow E$ that cosplits in $\mathcal{M}$ and such that $D \wedge_E D = E$.*

- (b) The canonical map $\text{Hom}_{\text{coalg}}(\varinjlim E_n, C) \rightarrow \text{Hom}_{\text{coalg}}(E_1, C)$ is surjective for every direct system of Hochschild extensions $\left((E_i)_{i \in \mathbb{N}}, \left(\eta_i^j\right)_{i,j \in \mathbb{N}}\right)$ which has direct limit $\varinjlim E_n$.
- (c) The canonical map $\text{Hom}_{\text{coalg}}(\varinjlim D^{\wedge_E^n}, C) \rightarrow \text{Hom}_{\text{coalg}}(D, C)$ is surjective for any coalgebra E in $\mathcal{M}$ and any subcoalgebra D of E such that $\left((D^{\wedge_E^i})_{i \in \mathbb{N}}, \left(\xi_i^j\right)_{i,j \in \mathbb{N}}\right)$ is a direct system of Hochschild extensions which has direct limit $\varinjlim D^{\wedge_E^n}$.
- (d) Any Hochschild extension of C is trivial.
- (e) $H^2(M, C) = 0$, for any $M \in {}^C\mathcal{M}^C$.

DEFINITION 4.12. Any coalgebra (C, Δ, ε) in $(\mathcal{M}, \otimes, \mathbf{1})$ satisfying one of the conditions of Theorem 4.11, is called *formally smooth*.

THEOREM 4.13. [AMS, Corollary 4.21] Let (C, Δ, ε) be a coalgebra in an abelian monoidal category $\mathcal{M}$. Then the following assertions are equivalent:

- (a) C is formally smooth.
- (b) Coker (Δ) is $\mathcal{I}$ -injective, where Δ is the comultiplication of C .

REMARK 4.14. We point out that, in view of Proposition 2.17, our definition of $D^{\wedge_E^n}$ and the one given in [AMS] agree.

THEOREM 4.15. Let (C, Δ, ε) be a formally smooth coalgebra in a cocomplete and complete abelian monoidal category $\mathcal{M}$ satisfying AB5, with left and right exact tensor functors. Assume that denumerable coproducts commute with $\otimes$. Let (M, ρ_M^r, ρ_M^l) be an $\mathcal{I}$ -injective C -bicomodule. Then the cotensor coalgebra $T_C^c(M)$ is formally smooth.

Proof. We will prove that any Hochschild extension of $T := T_C^c(M)$ is trivial. Let $\sigma : T \rightarrow B$ be a Hochschild extension of T . Since the canonical projection $p_0 : T \rightarrow C$ is a coalgebra homomorphism and C is formally smooth, by a) of Theorem 4.11, there exists a coalgebra homomorphism $f_C : B \rightarrow C$ such that $f_C \sigma = p_0$. Then B is a C -bicomodule via f_C . Moreover σ becomes a morphism of C -bicomodules. Since M is $\mathcal{I}$ -injective and the canonical projection $p_1 : T \rightarrow M$ is a morphism of C -bicomodules, then there is a morphism of C -bicomodules $f_M : B \rightarrow M$ such that $f_M \sigma = p_1$. Since, by Proposition 2.18, we have

$$\tilde{C}_T = \varinjlim C^{\wedge_T^n} = \varinjlim C^n(M) = T$$

since $\sigma_n : C^n \rightarrow T$ cosplits and since, by definition of Hochschild extension, $B = T^{\wedge_B^2}$ and σ cosplits, then by Theorem 3.11 applied to the case " α " = $i_0 : C \rightarrow T$ the canonical inclusion and " σ " = σ , we have $\tilde{C}_B = B$. Now we have

$$f_M \sigma i_0 = p_1 i_0 = 0.$$

Therefore we can apply Theorem 2.15 in the case when " C " = " D " = C , " M " = M , " E " = B and " δ " = σi_0 in order to obtain a unique coalgebra homomorphism $f : B \rightarrow T$ such that $p_0 f = f_C$ and $p_1 f = f_M$. Then we have

$$p_0 f \sigma = f_C \sigma = p_0, \quad \text{and} \quad p_1 f \sigma = f_M \sigma = p_1.$$

By Theorem 2.15 applied to $\delta = i_0 : C \rightarrow T$, the morphism Id_T is the unique coalgebra homomorphism such that $p_0 \text{Id}_T = p_0$ and $p_1 \text{Id}_T = p_1$. Therefore, we conclude that $f \sigma = \text{Id}_T$. $\square$

COROLLARY 4.16. Let (C, Δ, ε) be a coalgebra in a cocomplete and complete abelian monoidal category $\mathcal{M}$ satisfying AB5, with left and right exact tensor functors. Assume that denumerable coproducts commute with $\otimes$.

- a) If C is formally smooth, then
- a1) $T_C^c(C \otimes X \otimes C)$ is formally smooth, for any $X \in \mathcal{M}$. In particular $T_C^c(C^{\otimes n})$ is formally smooth, for any $n > 1$.
- a2) The cotensor coalgebra $T_C^c(\text{Coker}(\Delta))$ is formally smooth.
- b) If C is coseparable, then the cotensor coalgebra $T_C^c(M)$ is formally smooth, for any C -bicomodule M .

Proof. We will apply Theorem 4.15.

a1) By the dual of [AMS, Theo 1.16], all the objects of the form $C \otimes X \otimes C$, where $X \in \mathcal{M}$, are $\mathcal{I}$ -injective.

a2) By Theorem 4.13, since C is formally smooth, we have that $\text{Coker } \Delta$ is $\mathcal{I}$ -injective.

b) By Theorem 4.5, any C -bicomodule M is $\mathcal{I}$ -injective. $\square$

5. APPLICATIONS.

LEMMA 5.1. *Let R be a ring and let $\mathcal{M}$ be the category of right R -modules. Let $((X_i)_{i \in \mathbb{I}}, (\xi_i^j)_{i,j \in \mathbb{I}})$ be a direct system in $\mathcal{M}$, where, for $i \leq j$, $\xi_i^j : X_i \rightarrow X_j$, and let $(\xi_i : X_i \rightarrow C)_{i \in \mathbb{I}}$ be a compatible family of monomorphism in $\mathcal{M}$. Assume that $(C, (\xi_i)_{i \in \mathbb{I}}) = \varinjlim ((X_i)_{i \in \mathbb{I}}, (\xi_i)_{i \in \mathbb{I}})$. Then $C = \bigcup_{i \in \mathbb{I}} \text{Im}(\xi_i)$.*

Proof. Let $i \in \mathbb{I}$ and let $\tau_i : \text{Im}(\xi_i) \hookrightarrow C$, $\zeta_i : \text{Im}(\xi_i) \rightarrow \bigcup_{i \in \mathbb{I}} \text{Im}(\xi_i)$, $\tau : \bigcup_{i \in \mathbb{I}} \text{Im}(\xi_i) \rightarrow C$ be the canonical inclusions. We have that $\tau \zeta_i = \tau_i$, for any $i \in \mathbb{I}$. Note that, for $i \leq j$, we have $\text{Im}(\xi_i) \subseteq \text{Im}(\xi_j)$. Let $\alpha_i : X_i \rightarrow \text{Im}(\xi_i)$ be the corestriction to $\text{Im}(\xi_i)$ of ξ_i . Then α_i is an isomorphism. Note that $\tau_i \alpha_i = \xi_i$, for any $i \in \mathbb{I}$.

Since $(\zeta_i \alpha_i : X_i \rightarrow \bigcup_{i \in \mathbb{I}} \text{Im}(\xi_i))_{i \in \mathbb{I}}$ is a compatible family of morphisms, there exists a unique morphism $\lambda : C = \varinjlim ((X_i)_{i \in \mathbb{I}}, (\xi_i)_{i \in \mathbb{I}}) \rightarrow \bigcup_{i \in \mathbb{I}} \text{Im}(\xi_i)$ such that $\lambda \xi_i = \zeta_i \alpha_i$, for any $i \in \mathbb{I}$. Therefore we have

$$\tau \lambda \xi_i = \tau \zeta_i \alpha_i = \tau_i \alpha_i = \xi_i, \text{ for any } i \in \mathbb{I},$$

so that $\tau \lambda = \text{Id}_C$. Therefore τ is surjective, i.e. $C = \bigcup_{i \in \mathbb{I}} \text{Im}(\xi_i)$. $\square$

LEMMA 5.2. *Let $\delta : D \rightarrow C$ be a monomorphism which is a coalgebra homomorphism in the monoidal category $(\mathfrak{M}_K, \otimes_K, K)$ of vector spaces over a field K . Assume that $\tilde{D}_C = C$. Then $\text{Corad}(C) \subseteq \text{Im}(\delta)$.*

Proof. Denote by (L, p) the cokernel of δ in $\mathcal{M} = (\mathfrak{M}_K, \otimes_K, K)$. Recall that

$$(D^{\wedge_C^n}, \delta_n) := \ker(p^{\otimes n} \Delta_C^{n-1})$$

for any $n \in \mathbb{N} \setminus \{0\}$, where $\Delta^n : C \rightarrow C^{\otimes n+1}$ is the n^{th} iterated comultiplication of C . By Proposition 1.10, there are suitable morphisms $(\xi_i^j : D^{\wedge_C^i} \rightarrow D^{\wedge_C^j})_{i,j \in \mathbb{N}}$ such that $((D^{\wedge_C^i})_{i \in \mathbb{N}}, (\xi_i^j)_{i,j \in \mathbb{N}})$ is a direct system in $\mathcal{M}$ and $(\delta_n)_{n \in \mathbb{N}}$ is a compatible family. Moreover recall that $\tilde{D}_C = \varinjlim ((D^{\wedge_C^i})_{i \in \mathbb{N}}, (\xi_i)_{i \in \mathbb{N}})$ in $\mathcal{M}$. By Lemma 5.1, $C = \bigcup_{i \in \mathbb{N}} \text{Im}(\delta_i)$. Let $D_i := \text{Im}(\delta_{i+1})$ and note that

$$D_i = \{c \in C \mid p^{\otimes i+1} \Delta_C^i(c) = 0\}.$$

for any $i \geq 0$. Note that

$$(35) \quad \text{Im}(\delta) = D_0 \subseteq D_1 \subseteq \cdots \subseteq C.$$

By [Sw, Theorem 9.1.6, page 191], C is a filtered coalgebra, with coalgebra filtration given by (35) so that, by [Sw, Proposition 11.1.1, page 226], we have that $\text{Corad}(C) \subseteq D_0$. $\square$

THEOREM 5.3. *Let $\mathcal{M} = (\mathfrak{M}_K, \otimes_K, K)$ be the monoidal category of vector spaces over a field K , let E be a coalgebra in $\mathcal{M}$ and let M be a vector space. The following assertions are equivalent for a morphism $g : E \rightarrow M$ in $\mathcal{M}$:*

(i) $g(\text{Corad}(E)) = 0$.

(ii) *There is a monomorphism $\delta : D \rightarrow E$ which is a coalgebra homomorphism in $\mathcal{M}$ such that $\tilde{D}_E = E$ and $g\delta = 0$.*

Proof. (i) $\implies$ (ii) Choose $D := \text{Corad}(E)$ and let $\delta : D \rightarrow E$ be the canonical inclusion.

(ii) $\implies$ (i) By Lemma 5.2, we have $\text{Corad}(E) \subseteq \text{Im}(\delta)$. $\square$

REMARK 5.4. By means of Theorem 2.15 and Theorem 5.3, in the particular case when $\mathcal{M}$ is the category $(\mathfrak{M}_K, \otimes_K, K)$, of vector spaces over a field K , the universal property of our cotensor coalgebra is equivalent to Nichols's one.

5.5. A *monoidal functor* between two monoidal categories $(\mathcal{M}, \otimes, \mathbf{1}, a, l, r)$ and $(\mathcal{M}', \otimes, \mathbf{1}, a, l, r)$ is a triple (F, ϕ_0, ϕ_2) , where $F : \mathcal{M} \rightarrow \mathcal{M}'$ is a functor, $\phi_0 : \mathbf{1} \rightarrow F(\mathbf{1})$ is an isomorphism such that the diagrams

$$\begin{array}{ccc} \mathbf{1} \otimes F(U) & \xrightarrow{l_{F(U)}} & F(U) \\ \phi_0 \otimes F(U) \downarrow & & \uparrow F(l_U) \\ F(\mathbf{1}) \otimes F(U) & \xrightarrow{\phi_2(\mathbf{1}, U)} & F(\mathbf{1} \otimes U) \end{array} \quad \begin{array}{ccc} F(U) \otimes \mathbf{1} & \xrightarrow{r_{F(U)}} & F(U) \\ F(U) \otimes \phi_0 \downarrow & & \uparrow F(r_U) \\ F(U) \otimes F(\mathbf{1}) & \xrightarrow{\phi_2(U, \mathbf{1})} & F(U \otimes \mathbf{1}) \end{array}$$

are commutative, and $\phi_2(U, V) : F(U) \otimes F(V) \rightarrow F(U \otimes V)$ is a family of functorial isomorphisms such that the following diagram

$$\begin{array}{ccccc} (F(U) \otimes F(V)) \otimes F(W) & \xrightarrow{\phi_2(U, V) \otimes F(W)} & F(U \otimes V) \otimes F(W) & \xrightarrow{\phi_2(U \otimes V, W)} & F((U \otimes V) \otimes W) \\ \downarrow a_{F(U), F(V), F(W)} & & & & \downarrow F(a_{U, V, W}) \\ F(U) \otimes (F(V) \otimes F(W)) & \xrightarrow{F(U) \otimes \phi_2(V, W)} & F(U) \otimes F(V \otimes W) & \xrightarrow{\phi_2(U, V \otimes W)} & F(U \otimes (V \otimes W)) \end{array}$$

is commutative. A monoidal functor (F, ϕ_0, ϕ_2) is called *strict* if both ϕ_0 and ϕ_2 are identities.

The following Proposition states that the image of an (co)algebra through a monoidal functor carries a natural (co)algebra structure.

PROPOSITION 5.6. *Let $\mathcal{M}$ and $\mathcal{M}'$ be monoidal categories. Let (F, ϕ_0, ϕ_2) be a monoidal functor between the categories $\mathcal{M}$ and $\mathcal{M}'$. Then:*

1) *If (A, m, u) is an algebra in $\mathcal{M}$, then $(F(A), m_{F(A)}, u_{F(A)})$ is an algebra in $\mathcal{M}'$, where*

$$\begin{aligned} m_{F(A)} &:= F(A) \otimes F(A) \xrightarrow{\phi_2(A, A)} F(A \otimes A) \xrightarrow{F(m)} F(A) \\ u_{F(A)} &:= \mathbf{1}' \xrightarrow{\phi_0} F(\mathbf{1}) \xrightarrow{F(u)} F(A). \end{aligned}$$

Moreover $F(f) : F(A) \rightarrow F(B)$ is an algebra homomorphism in $\mathcal{M}'$ whenever $f : A \rightarrow B$ is an algebra homomorphism in $\mathcal{M}$. The converse also holds true if F is faithful.

2) *If (C, Δ, ε) is a coalgebra in $\mathcal{M}$, then $(F(C), \Delta_{F(C)}, \varepsilon_{F(C)})$ is a coalgebra in $\mathcal{M}'$, where*

$$\begin{aligned} \Delta_{F(C)} &:= F(C) \xrightarrow{F(\Delta)} F(C \otimes C) \xrightarrow{\phi_2^{-1}(C, C)} F(C) \otimes F(C) \\ \varepsilon_{F(C)} &:= F(C) \xrightarrow{F(\varepsilon)} F(\mathbf{1}) \xrightarrow{\phi_0^{-1}} \mathbf{1}'. \end{aligned}$$

Moreover $F(f) : F(C) \rightarrow F(D)$ is a coalgebra homomorphism in $\mathcal{M}'$ whenever $f : C \rightarrow D$ is a coalgebra homomorphism in $\mathcal{M}$. The converse also holds true if F is faithful.

Proof. follows directly from the definitions. □

Let K be a field. From now on $\mathfrak{M}_K$ will denote the category of vector spaces over K .

THEOREM 5.7. *Let $(\mathcal{M}, \otimes, \mathbf{1})$ be a complete and cocomplete abelian monoidal category. Let K be a field and let*

$$(F, \phi_0, \phi_2) : (\mathcal{M}, \otimes, \mathbf{1}) \rightarrow (\mathfrak{M}_K, \otimes_K, K)$$

be an additive monoidal functor. Assume that:

- (1) *F is faithful.*
- (2) *F preserves kernels and cokernels.*
- (3) *F preserves denumerable products and coproducts.*

Let C and E be coalgebras in $\mathcal{M}$. Let $f_C : E \rightarrow C$ be a coalgebra homomorphism and let $f_M : E \rightarrow M$ be a morphism of C -bicomodules, where E is a bicomodule via f_C . Assume that

$$F(f_M)(\text{Corad}(F(E))) = 0.$$

Then there is a unique coalgebra homomorphism $f : E \rightarrow T_C^c(M)$ such that $p_0 f = f_C$ and $p_1 f = f_M$, where $p_n : T_C^c(M) \rightarrow M^{\square n}$ denotes the canonical projection.

$$\begin{array}{ccc} T_C^c(M) & \xrightarrow{p_1} & M \\ p_0 \downarrow & \nearrow \scriptstyle f & \uparrow f_M \\ C & \xleftarrow{f_C} & E \end{array}$$

Proof. In order to simplify the computations, we will omit the isomorphisms ϕ_0, ϕ_2 .

For every object $X \in \mathcal{M}$, we will denote by X' the vector space $F(X)$. Also, for every morphism $f \in \mathcal{M}$, we will denote by f' the K -linear map $F(f)$.

Recall that E' and D' carry the coalgebra structures described in Proposition 5.6.

Since F preserves denumerable coproducts and cokernels, it preserves denumerable direct limits and finite coproducts so that

$$\begin{aligned} F[(T_C^c(M), (\sigma_n)_{n \in \mathbb{N}})] &= F\left[\varinjlim \left((C^n(M))_{n \in \mathbb{N}}, (\sigma_m^n)_{m, n \in \mathbb{N}}\right)\right] \\ &= \varinjlim \left((F[C^n(M)])_{n \in \mathbb{N}}, (F(\sigma_m^n))_{m, n \in \mathbb{N}}\right) \\ &= \varinjlim \left([(C')^n(M')]_{n \in \mathbb{N}}, (\sigma_m^n)_{m, n \in \mathbb{N}}\right). \end{aligned}$$

Therefore we get $(T_C^c(M))' = T_{C'}^c(M')$, that is the cotensor coalgebra in the category of vector spaces of the C' -bicomodule M' .

Now, by Proposition 5.6, $f'_C : E' \rightarrow C'$ is a coalgebra homomorphism and $f'_M : E' \rightarrow M'$ is a morphism of C' -bicomodules, where E' is a bicomodule via f'_C . By hypothesis, $f'_M(\text{Corad}(E')) = 0$.

By Theorem 5.3, there is a coalgebra D in $\mathfrak{M}_K$ and a monomorphism $\delta : D \rightarrow E'$ which is a coalgebra homomorphism in $\mathfrak{M}_K$ such that $\varinjlim D^{\wedge_{E'}^n} = E'$ and $f'_M \delta = 0$.

By Theorem 2.15, there exists a unique coalgebra homomorphism $g : E' \rightarrow T_{C'}^c(M')$ such that $p'_0 g = f'_C$ and $p'_1 g = f'_M$, where $p_n : T_{C'}^c(M') \rightarrow M'^{\square n}$ denotes the canonical projection:

$$\begin{array}{ccc} T_{C'}^c(M') & \xrightarrow{p'_1} & M' \\ p'_0 \downarrow & \nearrow \scriptstyle g & \uparrow f'_M \\ C' & \xleftarrow{f'_C} & E' \end{array}$$

We will prove that $g = f'$ for a suitable morphism $f : E \rightarrow T_C^c(M)$ in $\mathcal{M}$.

By construction (see (18)):

$$p'_k g = (f'_M)^{\square k} \overline{\Delta}_{E'}^{k-1} \text{ for any } k \in \mathbb{N}.$$

Let

$$q_k : \prod_{n \in \mathbb{N}} M^{\square n} \rightarrow M^{\square k}$$

be the canonical projection and let

$$\omega : T_C^c(M) = \bigoplus_{n \in \mathbb{N}} M^{\square n} \rightarrow \prod_{n \in \mathbb{N}} M^{\square n}$$

be the diagonal morphism of the $(p_k)_{k \in \mathbb{N}}$. This is uniquely defined by

$$q_k \omega = p_k, \text{ for every } k \in \mathbb{N}.$$

Since F preserves kernels and denumerable (co)products, it is clear that

$$\omega' : T_{C'}^c(M') = \bigoplus_{n \in \mathbb{N}} (M')^{\square n} \rightarrow \prod_{n \in \mathbb{N}} (M')^{\square n}$$

is a monomorphism. Since F is faithful, we get that ω is a monomorphism too. Let

$$\alpha : E \rightarrow \prod_{n \in \mathbb{N}} M^{\square n}$$

be the diagonal morphism of the $\left(f_M^{\square k} \circ \overline{\Delta}_E^{k-1}\right)_{k \in \mathbb{N}}$. This is uniquely defined by

$$q_k \alpha = f_M^{\square k} \circ \overline{\Delta}_E^{k-1}, \text{ for every } k \in \mathbb{N}.$$

We have that

$$\omega' g = \alpha'.$$

In fact

$$q'_k \omega' g = (q_k \omega)' g = p'_k g = (f'_M)^{\square k} \overline{\Delta}_{E'}^{k-1} = \left(f_M^{\square k} \circ \overline{\Delta}_E^{k-1}\right)' = (q_k \alpha)' = q'_k \alpha',$$

for every $k \in \mathbb{N}$.

Let $(\text{Coker}(\omega), \tau)$ be the cokernel of ω in $\mathcal{M}$. We have

$$(\tau \alpha)' = \tau' \alpha' = \tau' \omega' g = (\tau \omega)' g = 0.$$

Since F is faithful, we get $\tau \alpha = 0$. Since ω is a monomorphism, it is clear that $(T_C^c(M), \omega) = \text{Ker}(\tau)$. By the universal property of kernels, there exists a unique morphism $f : E \rightarrow T_C^c(M)$ in $\mathcal{M}$ such that

$$\omega f = \alpha.$$

We have

$$\omega' f' = (\omega f)' = \alpha' = \omega' g.$$

Since ω' is a monomorphism we get

$$f' = g.$$

By Proposition 5.6, we have that f is a coalgebra homomorphism.

It remains to prove that f is uniquely defined by the relations $p_0 f = f_C$ and $p_1 f = f_M$. Let h be another coalgebra homomorphism such that $p_0 h = f_C$ and $p_1 h = f_M$. Then

$$p'_0 h' = f'_C = p'_0 g \quad \text{and} \quad p'_1 h' = f'_M = p'_1 g.$$

By uniqueness of g , we get

$$h' = g = f'.$$

Since F is faithful we conclude that $h = f$. $\square$

5.8. Let H be a Hopf algebra over a field K . Consider the following examples of monoidal categories.

- The category ${}_H \mathfrak{M} = ({}_H \mathfrak{M}, \otimes_K, K)$, of all left modules over H . The tensor $V \otimes W$ of two left H -modules is an object in ${}_H \mathfrak{M}$ via the diagonal action; the unit is K regarded as a left H -module via ε_H .

- The category ${}_H \mathfrak{M}_H = ({}_H \mathfrak{M}_H, \otimes_K, K)$, of all two-sided modules over H . The tensor $V \otimes W$ of two H -bimodules carries, on both sides, the diagonal action; the unit is K regarded as a H -bimodule via ε_H .

- The category ${}^H \mathfrak{M} = ({}^H \mathfrak{M}, \otimes_K, K)$, of all left comodules over H . The tensor product $V \otimes W$ of two left H -comodules is an object in ${}^H \mathfrak{M}$ via the diagonal coaction; the unit is K regarded as a left H -comodule via the map $k \mapsto 1_H \otimes k$.

- The category ${}^H \mathfrak{M}^H = ({}^H \mathfrak{M}^H, \otimes_K, K)$ of all two-sided comodules over H . The tensor $V \otimes W$ of two H -bicomodules carries, on both sides, the diagonal coaction; the unit is K regarded as a H -bicomodule via the maps $k \mapsto 1_H \otimes k$ and $k \mapsto k \otimes 1_H$.

- The category ${}_H^H \mathcal{YD} = ({}_H^H \mathcal{YD}, \otimes_K, K)$ of left Yetter-Drinfeld modules over H . Recall that an object V in ${}_H^H \mathcal{YD}$ is a left H -module and a left H -comodule satisfying, for any $h \in H, v \in V$, the compatibility condition:

$$\sum (h_{(1)} v)_{<-1>} h_{(2)} \otimes (h_{(1)} v)_{<0>} = \sum h_{(1)} v_{<-1>} \otimes h_{(2)} v_{<0>}$$

or, equivalently,

$$\rho(hv) = \sum h_{(1)}v_{<-1>}S(h_{(3)}) \otimes h_{(2)}v_{<0>},$$

where $\Delta_H(h) = \sum h_{(1)} \otimes h_{(2)}$ and $\rho(v) = \sum v_{<-1>} \otimes v_{<0>}$ denote the comultiplication of H and the left H -comodule structure of V respectively (we used Sweedler notation).

The tensor product $V \otimes W$ of two Yetter-Drinfeld modules is an object in ${}^H_H\mathcal{YD}$ via the diagonal action and the codiagonal coaction; the unit in ${}^H_H\mathcal{YD}$ is K regarded as a left H -comodule via the map $x \mapsto 1_H \otimes x$ and as a left H -module via the counit ε_H .

Let $\mathcal{M}$ denote one of the categories above and let $F : \mathcal{M} \rightarrow \mathfrak{M}_K$ be the forgetful functor. Then Theorem 5.7 applies.

5.9. Let (H, ϕ) be a quasi-Hopf algebra over a field K . Then the category of right H -modules $\mathfrak{M}_H$ is a monoidal category. By [Maj, Example 9.1.4, page 422], the forgetful functor $\mathfrak{M}_H \rightarrow \mathfrak{M}_K$ is monoidal if and only if H is twisted-equivalent to an ordinary Hopf algebra. Therefore theorem 5.7 does not apply in general to $\mathfrak{M}_H$. Anyway Theorem 2.15 still holds. In fact, since H is in particular an ordinary K -algebra, it is clear that $\mathfrak{M}_H$ is a cocomplete and complete abelian monoidal category satisfying AB5.

REFERENCES

- [AG] N. Andruskiewitsch, M. Graña, *Braided Hopf algebras over non-abelian finite groups*. Colloquium on Operator Algebras and Quantum Groups (Spanish) (Vaquerías, 1997). Bol. Acad. Nac. Cienc. (Córdoba) 63 (1999), 45–78.
- [AS] N. Andruskiewitsch, H.-J. Schneider, *Pointed Hopf algebras*. New directions in Hopf algebras, 1–68, Math. Sci. Res. Inst. Publ., 43, Cambridge Univ. Press, Cambridge, 2002.
- [AMS] A. Ardizzoni, C. Menini and D. Ștefan, *Hochschild Cohomology and ‘Smoothness’ in Monoidal Categories*, submitted.
- [Ar1] A. Ardizzoni, *Separable Functors and Formal Smoothness*, preprint. arXiv math.QA/0407095.
- [Ar2] A. Ardizzoni, *The Category of Modules over a Monoidal Category: Abelian or not?*, Annali dell’Università di Ferrara, Sezione VII - Scienze Matematiche, in press.
- [CR] C. Cibils, M. Rosso, *Hopf quivers*. J. Algebra 254 (2002), no. 2, 241–251.
- [CQ] J. Cuntz and D. Quillen, *Algebra extensions and nonsingularity*, J. of AMS, **8** (1995), 251–289.
- [JLMS] P. Jara, D. Llena, L. Merino and D. Ștefan, *Hereditary and formally smooth coalgebras*, preprint.
- [Ka] C. Kassel, *Quantum groups*, Graduate Text in Mathematics **155**, Springer, 1995.
- [Maj] S. Majid, *Foundations of quantum group theory*, Cambridge University Press, 1995.
- [Mo] S. Montgomery, *Hopf Algebras and their actions on rings*, CMBS Regional Conference Series in Mathematics **82**, 1993.
- [Ni] Nichols, Warren D. Bialgebras of type one. Comm. Algebra 6 (1978), no. 15, 1521–1552.
- [Po] N. Popescu, *Abelian Categories with Application to Rings and Modules*, Academic Press, London & New York, (1973).
- [Ro] M. Rosso, *Quantum groups and quantum shuffles*. Invent. Math. 133 (1998), no. 2, 399–416.
- [St] B. Stenström, *Rings of quotients Die Grundlehren der Mathematischen Wissenschaften, Band 217. An introduction to methods of ring theory*. Springer-Verlag, New York-Heidelberg, 1975.
- [Sw] M. Sweedler, “Hopf Algebras”, Benjamin, New York, 1969.

A. ARDIZZONI.

UNIVERSITY OF FERRARA, DEPARTMENT OF MATHEMATICS, VIA MACHIAVELLI 35, FERRARA, I-44100, ITALY .
email: alessandro.ardizzoni@unife.it

URL: <http://www.unife.it/utenti/alessandro.ardizzoni>

C. MENINI.

UNIVERSITY OF FERRARA, DEPARTMENT OF MATHEMATICS, VIA MACHIAVELLI 35, FERRARA, I-44100, ITALY.
email: men@dns.unife.it

URL: <http://web.unife.it/utenti/claudia.menini>

D. ȘTEFAN.

UNIVERSITY OF BUCHAREST, FACULTY OF MATHEMATICS, STR. ACADEMIEI 14, BUCHAREST, RO-70109, ROMANIA.
email: dstefan@al.math.unibuc.ro